

White-matter relaxation time and myelin water fraction differences in young adults with autism

S. C. L. Deoni^{1†}, J. R. Zinkstok^{2*†}, E. Daly², C. Ecker², MRC AIMS Consortium[‡], S. C. R. Williams³
and D. G. M. Murphy^{2,4}

¹Advanced Baby Imaging Laboratory, School of Engineering, Brown University, Providence, RI, USA

²Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, King's College London, London, UK

³Department of Neuroimaging, Institute of Psychiatry, King's College London, London, UK

⁴The Sackler Institute for Translational Neurodevelopment, King's College London, London, UK

Background. Increasing evidence suggests that autism is associated with abnormal white-matter (WM) anatomy and impaired brain 'connectivity'. While myelin plays a critical role in synchronized brain communication, its aetiological role in autistic symptoms has only been *indirectly* addressed by WM volumetric, relaxometry and diffusion tensor imaging studies. A potentially more specific measure of myelin content, termed myelin water fraction (MWF), could provide improved sensitivity to myelin alteration in autism.

Method. We performed a cross-sectional imaging study that compared 14 individuals with autism and 14 age- and IQ-matched controls. T_1 relaxation times (T_1), T_2 relaxation times (T_2) and MWF values were compared between autistic subjects, diagnosed using the Autism Diagnostic Interview – Revised (ADI-R), with current symptoms assessed using the Autism Diagnostic Observation Schedule (ADOS) and typical healthy controls. Correlations between T_1 , T_2 and MWF values with clinical measures [ADI-R, ADOS, and the Autism Quotient (AQ)] were also assessed.

Results. Individuals with autism showed widespread WM T_1 and MWF differences compared to typical controls. Within autistic individuals, worse current social interaction skill as measured by the ADOS was related to reduced MWF although not T_1 . No significant differences or correlations with symptoms were observed with respect to T_2 .

Conclusions. Autistic individuals have significantly lower global MWF and higher T_1 , suggesting widespread alteration in tissue microstructure and biochemistry. Areas of difference, including thalamic projections, cerebellum and cingulum, have previously been implicated in the disorder; however, this is the first study to specifically indicate myelin alteration in these regions.

Received 6 December 2013; Revised 5 July 2014; Accepted 14 July 2014; First published online 11 August 2014

Key words: Autism, brain connectivity, brain development, human brain imaging, myelin, white matter.

Introduction

Autism spectrum disorder (ASD) is characterized by impairments in social interaction, communication and emotional processing, as well as by restrictive, stereotyped and repetitive behaviours and interests. There is increasing evidence that these symptoms have a biological basis. Individuals with autism show evidence of altered structural and functional 'connectivity' across large-scale brain systems (Koshino *et al.* 2005; Alexander

et al. 2007; Ben Bashat *et al.* 2007; Keller *et al.* 2007; Lee *et al.* 2007; Jones *et al.* 2010; Weng *et al.* 2010). This aberrant brain messaging may be related to, or mirrored by, altered brain development (Courchesne, 2004; Herbert *et al.* 2004; Courchesne *et al.* 2005) and differences in structural white and grey matter (Boddaert *et al.* 2004; McAlonan *et al.* 2005). A recurrent finding in children with ASD is that of increased overall brain volume, which has been suggested to result from differences in early brain development (Courchesne *et al.* 2001, 2007; Hazlett *et al.* 2005; Koshino *et al.* 2005; Wassink *et al.* 2007) and may be caused by differential effects driving white matter (WM) to be larger in the autistic brain. Specifically, those brain regions exhibiting the greatest volume increases correspond to later and prolonged myelinating pathways (Abell *et al.* 1999; Aylward *et al.* 2002; Carper & Courchesne, 2005). Further (Catani *et al.* 2008; Fields, 2008), these WM volume differences

* Address for correspondence: Dr J. Zinkstok, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, P023, De Crespigny Park, Denmark Hill, London SE5 8AF, UK.
(Email: janneke.zinkstok@kcl.ac.uk)

† These authors served as joint first authors.

‡ Members of the MRC AIMS Consortium are given in the Appendix.

persist into young adulthood and extend fronto-temporally and fronto-occipitally within connecting fibre tracts including the inferior and superior longitudinal fasciculi, occipitofrontal fasciculus, and the external capsule.

The first signs of autism usually present in the first 3 years of life, and this time-frame corresponds temporally with the most dynamic period of brain myelination. The elaboration of the myelin sheath along WM axons and development of the myelinated WM progresses rapidly over the first 2 years of life (Waber et al. 2007; Deoni et al. 2012), and is tempo-spatially associated with evolving cognitive and behavioural functioning (Nagy et al. 2004; Johnson et al. 2005). Myelin plays a critical role in establishing and maintaining congruent brain communication, and contributes substantially to WM volume. Histological evidence for abnormal myelination, myelin content, or myelin structure in the pathogenesis of ASD is derived from *ex vivo* post-mortem studies showing altered myelin composition with delayed compaction in the sheaths (Casanova, 2004, 2006; Palmen et al. 2004; Buxhoeveden et al. 2006). Insufficient compaction of the lipid myelin bilayers, or compositional irregularity, can diminish the conductive potential of the myelinated axon and result in uncoordinated information transfer. To date, however, *in vivo* investigations of myelin alteration in autism [and its possible relationship(s) to clinical symptomology] have been *indirect*. For example, findings of some magnetic resonance spectroscopy studies are consistent with altered phospholipid metabolism in prefrontal brain regions (Murphy et al. 2002; Carper & Courchesne, 2005). Studies of WM microstructure and micro-organization using diffusion tensor (DT)-MRI and relaxation time measurements (also termed relaxometry) have also provided indirect support for altered myelin content and structure. Cross-sectional DT-MRI studies have reported alterations in WM fractional anisotropy (FA), as well as mean and radial diffusivity (although considerable heterogeneity exists in the anatomical location of these differences), which may be related to altered myelin integrity alongside changes in fibre coherence and architecture. Developmentally, FA has also been reported to increase more slowly between 6 and 24 months of age in infants later diagnosed with autism (despite having increased FA at 6 months) (Wolff et al. 2012). Voxel-wise comparisons of T_2 and T_2^* relaxation times (Hendry et al. 2006) revealed an overall increase in cerebral WM T_2 in patients with autism. Regionally, T_2 was increased in associated WM of the bilateral primary sensory association areas in the parietal lobes, the visual association areas in the occipital lobes, and the WM underlying the supplementary motor areas (SMAs) in the frontal lobes. These observed

T_2 increase could reflect reduced myelin content, and correspondingly increased water content, in the ASD brain.

These studies represent important steps in elucidating microstructural abnormalities in the ASD brain. Although indirect, taken together, they support the suggestion that individuals with ASD have abnormalities in brain WM (and specifically myelin content) that may underpin some symptoms. However, to date, this hypothesis has not been directly tested *in vivo* due to the difficulty in specifically and quantitatively measuring myelin content (Beaulieu, 2002; Madler et al. 2008). For instance, while alterations in myelin content may influence DT-MRI and relaxation-time measurements, these are not specific or quantitative measures of myelin content as they also reflect other biophysical and biochemical features (e.g. water content, fibre architecture, density and coherence, and/or membrane permeability). Currently, the most robust approach to quantitatively estimating myelin content is through multi-component relaxation analysis (MCR) (Whittall et al. 1997). Within brain tissue, MCR aims to decompose the measured MR signal into contributions from two anatomically distinct water compartments: the slow relaxing intra- and extra-axonal water; and the faster relaxing water trapped between the myelin bilayers. Using MCR an estimate of the myelin-associated water pool's volume fraction, termed the myelin water fraction (MWF) is derived, which has been shown to correlate strongly with 'gold standard' histological estimates of myelin content (Beaulieu et al. 1998; Gareau et al. 2000; Webb et al. 2003; Laule et al. 2006, 2008). MCR has been used to investigate demyelinating disorders such as multiple sclerosis (MacKay et al. 2009; Kitzler et al. 2012; Kolind et al. 2012), as well as neurodevelopment in infants, toddlers and young children (Deoni et al. 2011, 2012).

We are the first to use a time-efficient MCR technique, termed mcDESPOT (multi-component-driven equilibrium single-pulse observation of T_1 and T_2) (Deoni et al. 2008), to compare MWF estimates in young adults with ASD and matched typically developing controls. We tested the main null hypothesis that people with autism exhibit no differences in MWF. We also tested the subsidiary null hypothesis that differences in brain MWF are not associated with variation in clinical symptoms. In addition to MWF, mcDESPOT also provides quantitative T_1 and T_2 relaxation-time estimates. We therefore tested the main and subsidiary null hypotheses using these measures to determine if prior studies of T_2 change in ASD (Hendry et al. 2006) were linked to alterations in myelin, or if they reflected changes in other tissue microstructure and biochemical features (i.e. iron content, water content, etc.).

Table 1. Summary of subject characteristics including age, and verbal IQ, performance IQ, and full-scale IQ scores (VIQ, PIQ, and FSIQ) as derived from the Wechsler Abbreviated Scale of Intelligence (WASI). All subjects were right-handed males. Means and standard errors of the mean (s.e.m.) are given; between-group differences in age, VIQ, PIQ, FSIQ, AQ, empathy quotient (EQ) and systemizing quotient (SQ) scores were calculated using *t* tests

	ASD individuals (N=14)	Controls (N=14)	Statistic	Significance (two-tailed)
Age, years (s.e.m.)	24.2 (1.21)	27.9 (1.55)	$t = -1.920$, $df = 28$	$p = 0.065$
FSIQ (s.e.m.)	107.1 (3.40)	109.3 (3.80)	$t = -0.419$, $df = 28$	$p = 0.679$
PIQ (s.e.m.)	105.7 (3.70)	110.7 (3.51)	$t = -1.099$, $df = 28$	$p = 0.281$
VIQ (s.e.m.)	105.7 (4.1)	106.6 (3.51)	$t = -0.351$, $df = 28$	$p = 0.728$
AQ (s.e.m.)	28.08 (2.77)	11.29 (1.33)	$t = 5.602$, $df = 28$	$p < 0.000$
EQ (s.e.m.)	24.62 (3.23)	46.23 (3.42)	$t = -4.60$, $df = 28$	$p < 0.000$
SQ (s.e.m.)	55.67 (6.97)	54.83 (5.29)	$t = -0.14$, $df = 28$	$p = 0.9$
ADOS (total)	11 (3.8)			
Communication (ADOS)	3.75 (0.44)			
Social interaction (ADOS)	7.67 (0.78)			
Repetitive behaviour (ADOS)	1.67 (0.052)			
ADI-R				
Communication (ADI-R)	12.86 (3.9)			
Social interaction (ADI-R)	17.64 (4.2)			
Repetitive behaviour (ADI-R)	4.7 (1.5)			

ADOS, Autism Diagnostic Observation Schedule; ADI-R, Autism Diagnostic Interview – Revised.

Methods and materials

Participants

Fourteen adult males with ASD and 14 healthy comparison male subjects (who did not differ significantly in age or IQ) were recruited into the study. All participants had a full-scale IQ <70 as measured by the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler, 1999). Participants with ASD were diagnosed according to ICD-10 research criteria (International Statistical Classification of Diseases and Health Related Problems – 10th revision; WHO, 2004) The initial clinical diagnosis was confirmed using the Autism Diagnostic Interview – Revised (ADI-R; Lord *et al.* 1994) and current symptoms were assessed using the Autism Diagnostic Observation Schedule (ADOS; Lord *et al.* 1989). All cases reached ADI-R algorithm cut-offs in the three domains of impaired social interaction, communication and repetitive behaviours and stereotyped patterns (see Table 1 for details). In addition, autistic traits were assessed in all participants using the Autism Quotient (AQ) – a self-report screening instrument that measures where an adult with normal intelligence lies on the continuum from autism to normality (Baron-Cohen *et al.* 2001; Woodbury-Smith *et al.* 2005).

Exclusion criteria for participants included: left handedness, history of intellectual disability (i.e. full-scale IQ <70), major psychiatric disorder (e.g. psychosis or the use of antipsychotic medication), head injury,

genetic disorder associated with autism (e.g. fragile X syndrome, tuberous sclerosis), or any other medical condition affecting brain function (e.g. epilepsy).

The study was approved by the National Research Ethics Committee, Suffolk. Informed written consent was obtained from all study participants.

Magnetic resonance imaging

Voxel-wise MWF maps were acquired of each participant using the mcDESPOT technique ($n = 46$), which involves the acquisition of series of T_1 -weighted SPOiled GRAdient recalled echo (SPGR, or spoiled FLASH) and T_1/T_2 -weighted balanced steady-state free precession (SSFP, FIESTA or TrueFISP) data over an incremented range of flip angles.

All imaging was performed on a GE Signa HDx 1.5 T clinical scanner equipped with an 8-channel head RF coil array. Whole-brain, sagittally oriented mcDESPOT data were acquired with a common $22\text{ cm} \times 22\text{ cm} \times 16\text{ cm}$ field of view (FOV) and $128 \times 128 \times 92$ imaging matrix. To reduce acquisition time, data were acquired with 3/4 partial Fourier acquisition. Sequence-specific acquisition parameters were as follows:

SPGR. Echo time (TE)/repetition time (TR) = 2.5 ms/5.3 ms, flip angles (α) = 3°, 4°, 5°, 6°, 7°, 9°, 12° and 17°, receiver bandwidth (BW) = $\pm 22.3\text{ kHz}$.

SSFP. TE/TR = 1.6 ms/3.2 ms, α = 12°, 16°, 21°, 27°, 33°, 40°, 5°1 and 68°, BW = $\pm 50\text{ kHz}$.

The SSFP data were acquired with two phase-cycling increments (0° and 180°), allowing correction for main magnetic field (B_0) off-resonance effects (Deoni, 2009). An inversion prepared (IR-)SPGR image was also acquired (same FOV, $128 \times 64 \times 46$ acquisition matrix, TE/TR/inversion time/ $\alpha = 2.5$ ms/5.3 ms/350 ms/ 5° ; and BW = ± 22.3 kHz) to correct for flip angle (B_1) inhomogeneity (Deoni, 2007).

Data analysis

T_1 , T_2 and MWF calculation

Following acquisition, each participant's data were linearly co-registered to correct for subtle intra-session motion (Jenkinson et al. 2002), non-parenchyma signal was removed (Smith, 2002), and voxel-wise T_1 , T_2 and MWF estimates were calculated as described in Deoni et al. (2013), providing 3-dimensional 'maps' of these parameters. These maps from each participant were then nonlinearly co-registered to custom T_1 -weighted template constructed from a subsample of six healthy and six ASD participants. This template was created from the high flip angle T_1 -weighted SPGR image of each included participant using symmetric diffeomorphic normalization (SyN; Avants et al. 2008) as implemented in the ANTs package, and a cross-correlation similarity measure (<http://picsl.upenn.edu/ANTS>), using the `buildtemplateparallel.sh` script distributed with the ANTs package (Avants et al. 2010). Subsequent registration of each study participant's MWF, T_1 and T_2 maps to the common template was accomplished by first nonlinearly co-registering each participant's high flip angle T_1 -weighted SPGR image to the common template and then applying the calculated transformation matrix to the corresponding T_1 , T_2 and MWF maps.

Group-wise comparisons of T_1 , T_2 and MWF

Group-wise comparison of T_1 , T_2 and MWF between the ASD and matched typically developing controls was performed via voxel-wise unpaired two-tailed t tests, with cluster-based correction for multiple comparisons and false discovery. A 3 mm full-width-at-half-maximum Gaussian kernel was used to smooth the T_1 , T_2 and MWF data, and non-parametric permutation testing used to perform the group comparison (performed using the *randomize* tool included in the FMRIB Software Library; <http://www.fmrib.ox.ac.uk/fsl/>). Cluster-based correction was performed using a cluster threshold of 2.5. Significance was defined as $p < 0.05$, cluster corrected.

Correlations between T_1 , T_2 and MWF and symptom measures

To examine the subsidiary hypothesis that altered T_1 , T_2 and/or MWF are associated with autism spectrum

traits, voxel-wise correlation analysis was performed in ASD individuals between T_1 , T_2 and MWF, and total ADOS scores, as well as the communication, social and repetitive domains subscores; ADI-R communication, social, repetitive, and developmental domain subscores; and total AQ scores.

Similar correlation analysis was also performed between T_1 , T_2 and MWF, and total AQ score within the typically developing control group only. Finally, correlations between T_1 , T_2 and MWF, and total AQ score were investigated in the combined ASD + typically developing control group.

All correlation analysis was performed via the *randomize* tool using non-parametric testing, with cluster-based correction for multiple comparisons (threshold = 2.5) and statistical significance was defined as $p < 0.05$, cluster corrected.

Results

Demographic data (Table 1)

A summary of the healthy and ASD cohorts, including mean age, verbal IQ, performance IQ, full-scale IQ, AQ, as well as ADOS and ADI-R total and subscore measures are provided in Table 1.

The groups did not differ significantly in mean age, full-scale, verbal, or performance IQ scores. As expected, ASD subjects had significantly higher AQ scores relative to the healthy participants.

Mean cerebral and cerebellar volumes and total WM volumes did not differ significantly between individuals with autism and healthy controls (Table 2).

Group-wise comparisons of T_1 , T_2 and MWF (Fig. 1)

Widespread differences in MWF (all significantly reduced in individuals with ASD compared to controls) and T_1 (all significantly increased in individuals with ASD compared to controls) were found. No significant differences in T_2 were observed. From the MWF results, significantly ($p < 0.05$) reduced MWF was observed in ASD individuals bilaterally within the cerebellum, thalamus, internal capsule, caudate nuclei, temporal and occipital WM, the SMA and pre-SMA, and cingulum; and right frontal WM. Significantly ($p < 0.05$) increased T_1 was observed in ASD individuals bilaterally within the cerebellum, thalamus, and internal capsule; and in right temporal and occipital WM.

Correlations between T_1 , T_2 and MWF and symptom measures (Figs 2 and 3)

Within individuals with ASD, there was a significant ($p < 0.05$) negative correlation between: (1) total ADOS score and MWF (i.e. lower MWF was associated with

Table 2. Summary of mean volumes (ml) with standard deviations (s.d.) and between-group differences calculated using t tests

	Patients (N=14)	Controls (N=14)	Significance (two-tailed)
Cerebrum (s.d.)	569.10 (41.34)	556.21 (32.25)	$p=0.37$
Left cerebellum (s.d.)	17.12 (3.75)	16.05 (2.63)	$p=0.39$
Right cerebellum (s.d.)	16.22 (3.74)	15.24 (2.16)	$p=0.41$
Total white matter (s.d.)	602.44 (46.31)	587.50 (34.66)	$p=0.34$




Fig. 1. Groupwise myelin water fraction (MWF), T_1 and T_2 comparisons between individuals with autism spectrum disorder (ASD) and healthy controls. Individuals with ASD show widespread significant ($p < 0.05$) reductions in MWF, with associated increased T_1 . No significant differences were observed in T_2 .

more abnormal current behaviour) bilaterally in the thalamus and caudate; and in the right frontal lobe and external capsule (Fig. 2); (2) negative correlation between ADOS social interaction subscore and MWF bilaterally in the thalamus; and in the right temporal and frontal lobes and cingulum (Fig. 2); and (3) negative correlation between AQ score and MWF in the right

cerebellum, occipital lobe and superior corona radiata (Fig. 2). We found no significant associations between MWF and ADI or any other subscales of the ADOS or ADI. We found no significant associations between T_1 or T_2 and any ADI, ADOS or AQ measure.

When we examined the dimensional relationship between autistic traits and MWF across all individuals




Fig. 2. Areas of significant ($p < 0.05$) negative correlation between myelin water fraction (MWF) and Autism Diagnostic Observation Schedule (ADOS) total score; MWF and ADOS social interaction subscore; and MWF and Autism Quotient (AQ) score with the autism spectrum disorder (ASD) subjects only. In these regions, reduced MWF is associated with more abnormal behaviour.

(i.e. combining ASD and typically developing participants), we found a significant ($p < 0.05$) *negative* correlation between MWF and AQ score bilaterally in the thalamus and cerebellum (Fig. 3). No significant correlations were found between either T_1 or T_2 and AQ score in the combined group.

Discussion

In this work, we have examined if individuals with ASD have significant differences in brain myelin content, or MWF, and if these differences may, in part, explain prior findings of T_2 relaxation time differences in ASD. This is the first study to use a whole-brain

multicomponent relaxometry approach to more specifically interrogate myelin content differences in ASD. While the number of participants in the study was small (14 in each group), the individuals with autism were free of many potential confounds that affect brain structure and function (e.g. epilepsy and/or intellectual disability). This homogeneous sample provided sufficient power to detect highly significant group differences.

Our results show that individuals with autism have widespread MWF reductions in brain regions previously implicated in ASD. Increased T_1 was also found in areas exhibiting lower MWF measures; however, the extent of T_1 differences was smaller than




Fig. 3. Areas of significant ($p < 0.05$) negative correlation between myelin water fraction (MWF) and Autism Quotient (AQ) scores in the combined autism spectrum disorder (ASD) plus control subjects. In these areas, reduced MWF is associated with higher AQ scores.

MWF differences. No group T_2 differences were found. Spatial inconsistencies in MWF, T_1 and T_2 changes reaffirm that these measures reflect different aspects of tissue microstructure. While both T_1 and T_2 are affected by changes in water, macromolecule, lipid and protein content, T_2 is also sensitive to changes in iron and other paramagnetic material content. MWF is believed to be more specific to changes in lipid myelin content. Our observation of T_1 and MWF differences suggests that myelin lipid and water content are altered in ASD (Alexander *et al.* 2011).

Within the ASD group, a significant negative correlation was found between MWF and the social domain of the ADOS in regions that are known to be involved in social processing. Previous DT-MRI studies of structural connectivity have, likewise, shown abnormal WM microstructure in pathways through, or connecting, these regions (Cheng *et al.* 2010). The absence of a similar finding with respect to ADI-R scores is likely reflective of the ADOS measuring current behaviour, whereas the ADI-R is mainly a measure of past behaviour.

Our findings of widespread disturbances in WM myelin content are consistent with, and add to, prior reports of WM alterations in autism. DT-MRI studies have shown reduced FA (an indicator of microstructural coherence) in the corpus callosum, internal capsule, and other WM regions (e.g. see Courchesne, 2004; Herbert *et al.* 2004; Courchesne *et al.* 2005; Jones *et al.* 2010, among others). Voxel-based morphometric analyses have reported widespread differences in WM volume (or density) (Ben Bashat *et al.* 2007; Lee *et al.* 2007). Unfortunately, relating these observed FA or morphometric findings to specific WM disruptions (i.e. myelin loss, reduced axonal size or density, etc.) has been challenging. Alterations in FA, for example, can reflect changes in myelin, fibre coherence, density, etc. Likewise, alterations in WM density can reflect

fibre density and water content (Beaulieu, 2002), as well as inconsistencies in MR tissue contrast. Thus, while these prior studies have consistently demonstrated altered WM in autism, they have been unable to elucidate the particular mechanism(s) responsible. Our results are the first to specifically investigate the role of myelin in autism, and to link observed changes in MWF to autistic traits and symptoms.

The development of myelin (myelination) plays a critical role in brain development. The formation of efficient information pathways throughout the brain is essential for normal function, cognition and behaviour. Our study has revealed widespread myelin alteration through MWF reduction in the brain of adults with autism. However, a limitation of our cross-sectional study design is that we cannot state when these deficits first arose, or whether they are secondary to other pathological processes. For example, prematurity, low birth weight, and ischaemic injury at birth have all been associated with WM microstructural changes, as assessed with DT-MRI, in later life (Skranes *et al.* 2007; Allen, 2008; Constable *et al.* 2008; Rose *et al.* 2008). However, to the best of our knowledge, no *in vivo* study has yet investigated brain myelination in children or adults who had these birth difficulties and, thus, how myelination is affected. To investigate these crucial questions, longitudinal studies of myelination in typically developing children and children at risk for autism, controlling for prematurity, etc., are required.

Associations between myelin development and the genetic and epigenetic factors contributing to autism also warrant further investigation. For example, knowledge of the timing, location and degree to which gene expression is disrupted in the pathway to the development of ASD is crucial for developing therapeutic strategies (Geschwind & Levitt, 2007). To date, however, few human studies have investigated

myelin-associated genes, including myelin basic protein (MBP), proteolipid protein (PLP) or myelin oligodendrocyte glycoprotein (MOG), in ASD. Animal studies, however, have shown the expression of genes associated with MBP, MOG and PLP are altered following prenatal viral infection, which some have reported as being associated with the development of autism (Fatemi *et al.* 2009). Additionally, the *SLC25A12* (solute carrier family 25 member 12) gene, identified by some as an autism susceptibility gene (Ramos *et al.* 2004), has recently been associated with global hypomyelination (Wibom *et al.* 2009). Although preliminary, these reports suggest the need for further combined imaging and genetic studies of myelination in ASD.

A further question not addressed in our work is whether the detected MWF abnormalities are *causative* of observed clinical deficits, or if they are symptoms of other pathological processes. For example, myelination of the brain in early infancy is believed to follow a pattern that spatially and temporally corresponds to the developing neuronal systems (Paus *et al.* 2001; Durston & Casey, 2006). However, it is not yet known if, for example, abnormal myelination of language pathways results in poor language performance, or if inadequate learning of language results in under-myelination of those neural pathways. Again, understanding of these relationships, and how they are affected in ASD, necessitates longitudinal studies of young children with autism or infants at risk of developing the disorder.

Our results, however, do show that autistic traits are associated with lower MWF measures within relevant WM pathways and are consistent with prior reports of microstructural abnormality and hypothesized aberrant connectivity. Myelin plays a critical role in facilitating coordinated information transfer throughout the brain (as evidenced by the loss of function and cognition observed in demyelinating disorders such as multiple sclerosis). Altered myelin, therefore, is likely associated with reduced connectivity. Our results are consistent with the current hypothesis that neural disconnectivity underpins ASD (Belmonte *et al.* 2004; Alexander *et al.* 2007; Hughes, 2007; Kleinhans *et al.* 2008), as supported by structural imaging studies (Barnea-Goraly *et al.* 2004; Ben Bashat *et al.* 2007); functional imaging studies (Castelli *et al.* 2002; Villalobos *et al.* 2005; Weng *et al.* 2010); and electroencephalography investigations (Grice *et al.* 2001; Brown *et al.* 2005). In each of these prior studies, abnormal connectivity was observed in frontal and temporal regions, consistent with our findings of lower myelin content in these areas.

In conclusion, the results of our study demonstrate for the first time that adults with ASD have highly

significant (widespread) differences in myelin content (as measured by MWF) compared to age- and IQ-matched controls; and that differences in myelin content in some brain regions are related to clinical symptoms and autistic traits.

Appendix

The MRC AIMS Consortium is a collaboration of autism research centres in the UK including the Institute of Psychiatry, London, The Autism Research Centre, University of Cambridge, and the Autism Research Group, University of Oxford. It is funded by the Medical Research Council (MRC) UK and headed by the Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry. The Consortium members are (in alphabetical order): Bailey AJ, Baron-Cohen S, Bolton PF, Bullmore ET, Carrington S, Chakrabarti B, Daly EM, Deoni SC, Ecker C, Happe F, Henty J, Jezzard P, Johnston P, Jones DK, Lombardo M, Madden A, Mullins D, Murphy CM, Murphy DG, Pasco G, Sadek S, Spain D, Steward R, Suckling J, Wheelwright S and Williams SC.

Acknowledgements

This work was supported by the Autism Imaging Multicentre Study Consortium, Medical Research Council UK Grant G0400061, and by European Autism Interventions – A Multicentre Study for Developing New Medications (EU-AIMS), which receives support from the Innovative Medicines Initiative Joint Undertaking under grant agreement no. 115300; and by a Medical Research Council fellowship awarded to S.D. (G0800298). We thank the National Institute for Health Research Biomedical Research Centre for Mental Health, and the Dr Mortimer and Theresa Sackler Foundation for their financial support.

Declaration of Interest

None.

References

- Abell F, Krams M, Ashburner J, Passingham R, Friston K, Frackowiak R, Happe F, Frith C, Frith U (1999). The neuroanatomy of autism: a voxel-based whole brain analysis of structural scans. *Neuroreport* 10, 1647–1651.
- Alexander AL, Hurley SA, Samsonov AA, Adluru N, Hosseinbor AP, Mossahebi P, Tromp do PM, Zakszewski E, Field AS (2011). Characterization of cerebral white matter properties using quantitative magnetic resonance imaging stains. *Brain Connectivity* 6, 423–446.

- Alexander AL, Lee JE, Lazar M, Boudos R, DuBray MB, Oakes TR, Miller JN, Lu J, Jeong EK, McMahon WM, Bigler ED, Lainhart JE (2007). Diffusion tensor imaging of the corpus callosum in Autism. *Neuroimage* **34**, 61–73.
- Allen MC (2008). Neurodevelopmental outcomes of preterm infants. *Current Opinion in Neurology* **21**, 123–128.
- Avants BB, Epstein CL, Grossman M, Gee JC (2008). Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis* **12**, 26–41.
- Avants BB, Yushkevich P, Pluta J, Minkoff D, Korczykowski M, Detre J, Gee JC (2010). The optimal template effect in hippocampus studies of diseased populations. *Neuroimage* **49**, 2457–2466.
- Aylward EH, Minshew NJ, Field K, Sparks BF, Singh N (2002). Effects of age on brain volume and head circumference in autism. *Neurology* **59**, 175–183.
- Barnea-Goraly N, Kwon H, Menon V, Eliez S, Lotspeich L, Reiss AL (2004). White matter structure in autism: preliminary evidence from diffusion tensor imaging. *Biological Psychiatry* **55**, 323–326.
- Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I (2001). The Reading the Mind in the Eyes Test revised version: a study with normal adults, and adults with Asperger syndrome or high-functioning autism. *Journal of Child Psychology and Psychiatry* **42**, 241–251.
- Beaulieu C (2002). The basis of anisotropic water diffusion in the nervous system – a technical review. *NMR in Biomedicine* **15**, 435–455.
- Beaulieu C, Fenrich FR, Allen PS (1998). Multicomponent water proton transverse relaxation and T₂-discriminated water diffusion in myelinated and nonmyelinated nerve. *Magnetic Resonance Imaging* **16**, 1201–1210.
- Belmonte MK, Allen G, Beckel-Mitchener A, Boulanger LM, Carper RA, Webb SJ (2004). Autism and abnormal development of brain connectivity. *Journal of Neuroscience* **24**, 9228–9231.
- Ben Bashat D, Kronfeld-Duenias V, Zachor DA, Ekstein PM, Hendler T, Tarrasch R, Even A, Levy Y, Ben Sira L (2007). Accelerated maturation of white matter in young children with autism: a high b value DWI study. *Neuroimage* **37**, 40–47.
- Boddaert N, Chabane N, Gervais H, Good CD, Bourgeois M, Plumet MH, Barthelemy C, Mouren MC, Artiges E, Samson Y, Brunelle F, Frackowiak RS, Zilbovicius M (2004). Superior temporal sulcus anatomical abnormalities in childhood autism: a voxel-based morphometry MRI study. *Neuroimage* **23**, 364–369.
- Brown C, Gruber T, Boucher J, Rippon G, Brock J (2005). Gamma abnormalities during perception of illusory figures in autism. *Cortex* **41**, 364–376.
- Buxhoeveden DP, Semendeferi K, Buckwalter J, Schenker N, Switzer R, Courchesne E (2006). Reduced minicolumns in the frontal cortex of patients with autism. *Neuropathology and Applied Neurobiology* **32**, 483–491.
- Carper RA, Courchesne E (2005). Localized enlargement of the frontal cortex in early autism. *Biological Psychiatry* **57**, 126–133.
- Casanova MF (2004). White matter volume increase and minicolumns in autism. *Annals of Neurology* **56**, 453; author reply 454.
- Casanova MF (2006). Neuropathological and genetic findings in autism: the significance of a putative minicolumnopathy. *Neuroscientist* **12**, 435–441.
- Castelli F, Frith C, Happe F, Frith U (2002). Autism, Asperger syndrome and brain mechanisms for the attribution of mental states to animated shapes. *Brain* **125**, 1839–1849.
- Catani M, Jones DK, Daly E, Embiricos N, Deeley Q, Pugliese L, Curran S, Robertson D, Murphy DG (2008). Altered cerebellar feedback projections in Asperger syndrome. *Neuroimage* **41**, 1184–1191.
- Cheng Y, Chou KH, Chen IY, Fan YT, Decety J, Lin CP (2010). Atypical development of white matter microstructure in adolescents with autism spectrum disorders. *Neuroimage* **50**, 873–882.
- Constable RT, Ment LR, Vohr BR, Kesler SR, Fulbright RK, Lacadie C, Delancy S, Katz KH, Schneider KC, Schafer RJ, Makuch RW, Reiss AR (2008). Prematurely born children demonstrate white matter microstructural differences at 12 years of age, relative to term control subjects: an investigation of group and gender effects. *Pediatrics* **121**, 306–316.
- Courchesne E (2004). Brain development in autism: early overgrowth followed by premature arrest of growth. *Mental Retardation and Developmental Disabilities Research Reviews* **10**, 106–111.
- Courchesne E, Karns CM, Davis HR, Ziccardi R, Carper RA, Tigue ZD, Chisum HJ, Moses P, Pierce K, Lord C, Lincoln AJ, Pizzo S, Schreibman L, Haas RH, Akshoomoff NA, Courchesne RY (2001). Unusual brain growth patterns in early life in patients with autistic disorder: an MRI study. *Neurology* **57**, 245–254.
- Courchesne E, Pierce K, Schumann CM, Redcay E, Buckwalter JA, Kennedy DP, Morgan J (2007). Mapping early brain development in autism. *Neuron* **56**, 399–413.
- Courchesne E, Redcay E, Morgan JT, Kennedy DP (2005). Autism at the beginning: microstructural and growth abnormalities underlying the cognitive and behavioral phenotype of autism. *Development and Psychopathology* **17**, 577–597.
- Deoni SC (2007). High-resolution T₁ mapping of the brain at 3T with driven equilibrium single pulse observation of T₁ with high-speed incorporation of RF field inhomogeneities (DESPOT1-HIFI). *Journal of Magnetic Resonance Imaging* **26**, 1106–1111.
- Deoni SC (2009). Transverse relaxation time (T₂) mapping in the brain with off-resonance correction using phase-cycled steady-state free precession imaging. *Journal of Magnetic Resonance Imaging* **30**, 411–417.
- Deoni SC, Dean DC, O’Muirheartaigh J, Dirks H, Jerskey BA (2012). Investigating white matter development in infancy and early childhood using myelin water fraction and relaxation time mapping. *Neuroimage* **63**, 1038–1053.
- Deoni SC, Matthews L, Kolind SH (2013). One component? Two components? Three? The effect of including a non-exchanging free water component in multicomponent

- driven equilibrium single pulse observation of T_2 and T_2 . *Magnetic Resonance Medicine* **70**, 147–154.
- Deoni SC, Mercure E, Blasi A, Gasston D, Thomson A, Johnson M, Williams SC, Murphy DG** (2011). Mapping infant brain myelination with magnetic resonance imaging. *Journal of Neuroscience* **31**, 784–791.
- Deoni SC, Rutt BK, Arun T, Pierpaoli C, Jones DK** (2008). Gleaning multicomponent T_1 and T_2 information from steady-state imaging data. *Magnetic Resonance Medicine* **60**, 1372–1387.
- Durston S, Casey BJ** (2006). What have we learned about cognitive development from neuroimaging? *Neuropsychologia* **44**, 2149–2157.
- Fatemi SH, Folsom TD, Reutiman TJ, Abu-Odeh D, Mori S, Huang H, Oishi K** (2009). Abnormal expression of myelination genes and alterations in white matter fractional anisotropy following prenatal viral influenza infection at E16 in mice. *Schizophrenia Research* **112**, 46–53.
- Fields RD** (2008). White matter in learning, cognition and psychiatric disorders. *Trends in Neuroscience* **31**, 361–370.
- Gareau PJ, Rutt BK, Karlik SJ, Mitchell JR** (2000). Magnetization transfer and multicomponent T_2 relaxation measurements with histopathologic correlation in an experimental model of MS. *Journal of Magnetic Resonance Imaging* **11**, 586–595.
- Geschwind DH, Levitt P** (2007). Autism spectrum disorders: developmental disconnection syndromes. *Current Opinion in Neurobiology* **17**, 103–111.
- Grice SJ, Spratling MW, Karmiloff-Smith A, Halit H, Csibra G, de Haan M, Johnson MH** (2001). Disordered visual processing and oscillatory brain activity in autism and Williams syndrome. *Neuroreport* **12**, 2697–2700.
- Hazlett HC, Poe M, Gerig G, Smith RG, Provenzale J, Ross A, Gilmore J, Piven J** (2005). Magnetic resonance imaging and head circumference study of brain size in autism: birth through age 2 years. *Archives of General Psychiatry* **62**, 1366–1376.
- Hendry J, DeVito T, Gelman N, Densmore M, Rajakumar N, Pavlosky W, Williamson PC, Thompson PM, Drost DJ, Nicolson R** (2006). White matter abnormalities in autism detected through transverse relaxation time imaging. *Neuroimage* **29**, 1049–1057.
- Herbert MR, Ziegler DA, Makris N, Filipek PA, Kemper TL, Normandin JJ, Sanders HA, Kennedy DN, Caviness VS Jr.** (2004). Localization of white matter volume increase in autism and developmental language disorder. *Annals of Neurology* **55**, 530–540.
- Hughes JR** (2007). Autism: the first firm finding=underconnectivity? *Epilepsy and Behaviour* **11**, 20–24.
- Jenkinson M, Bannister P, Brady M, Smith S** (2002). Improved optimization for the robust and accurate linear registration and motion correction of brain images. *Neuroimage* **17**, 825–841.
- Johnson MH, Griffin R, Csibra G, Halit H, Farroni T, de Haan M, Tucker LA, Baron-Cohen S, Richards J** (2005). The emergence of the social brain network: evidence from typical and atypical development. *Development and Psychopathology* **17**, 599–619.
- Jones TB, Bandettini PA, Kenworthy L, Case LK, Milleville SC, Martin A, Birn RM** (2010). Sources of group differences in functional connectivity: an investigation applied to autism spectrum disorder. *Neuroimage* **49**, 401–414.
- Keller TA, Kana RK, Just MA** (2007). A developmental study of the structural integrity of white matter in autism. *Neuroreport* **18**, 23–27.
- Kitzler HH, Su J, Zeineh M, Harper-Little C, Leung A, Kremenchutzky M, Deoni SC, Rutt BK** (2012). Deficient MWF mapping in multiple sclerosis using 3D whole-brain multi-component relaxation MRI. *Neuroimage* **59**, 2670–2677.
- Kleinhans NM, Richards T, Sterling L, Stegbauer KC, Mahurin R, Johnson LC, Greenson J, Dawson G, Aylward E** (2008). Abnormal functional connectivity in autism spectrum disorders during face processing. *Brain* **131**, 1000–1012.
- Kolind S, Matthews L, Johansen-Berg H, Leite MI, Williams SC, Deoni S, Palace J** (2012). Myelin water imaging reflects clinical variability in multiple sclerosis. *Neuroimage* **60**, 263–270.
- Koshino H, Carpenter PA, Minshew NJ, Cherkassky VL, Keller TA, Just MA** (2005). Functional connectivity in an fMRI working memory task in high-functioning autism. *Neuroimage* **24**, 810–821.
- Laule C, Kozlowski P, Leung E, Li DK, Mackay AL, Moore GR** (2008). Myelin water imaging of multiple sclerosis at 7T: correlations with histopathology. *Neuroimage* **40**, 1575–1580.
- Laule C, Leung E, Lis DK, Traboulsee AL, Paty DW, MacKay AL, Moore GR** (2006). Myelin water imaging in multiple sclerosis: quantitative correlations with histopathology. *Multiple Sclerosis* **12**, 747–753.
- Lee JE, Bigler ED, Alexander AL, Lazar M, DuBray MB, Chung MK, Johnson M, Morgan J, Miller JN, McMahon WM, Lu J, Jeong EK, Lainhart JE** (2007). Diffusion tensor imaging of white matter in the superior temporal gyrus and temporal stem in autism. *Neuroscience Letters* **424**, 127–132.
- Lord C, Rutter M, Goode S, Heemsbergen J, Jordan H, Mawhood L, Schopler E** (1989). Autism diagnostic observation schedule: a standardized observation of communicative and social behavior. *Journal of Autism and Developmental Disorders* **19**, 185–212.
- Lord C, Rutter M, Le Couteur A** (1994). Autism Diagnostic Interview – Revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders* **24**, 659–685.
- MacKay AL, Vavasour IM, Rauscher A, Kolind SH, Madler B, Moore GR, Traboulsee AL, Li DK, Laule C** (2009). MR relaxation in multiple sclerosis. *Neuroimaging Clinics of North America* **19**, 1–26.
- Madler B, Drabycz SA, Kolind SH, Whittall KP, MacKay AL** (2008). Is diffusion anisotropy an accurate monitor of myelination? Correlation of multicomponent T_2 relaxation and diffusion tensor anisotropy in human brain. *Magnetic Resonance Imaging* **26**, 874–888.

- McAlonan GM, Cheung V, Cheung C, Suckling J, Lam GY, Tai KS, Yip L, Murphy DG, Chua SE (2005). Mapping the brain in autism. A voxel-based MRI study of volumetric differences and intercorrelations in autism. *Brain* **128**, 268–276.
- Murphy DG, Critchley HD, Schmitz N, McAlonan G, Van Amelsvoort T, Robertson D, Daly E, Rowe A, Russell A, Simmons A, Murphy KC, Howlin P (2002). Asperger syndrome: a proton magnetic resonance spectroscopy study of brain. *Archives of General Psychiatry* **59**, 885–891.
- Nagy Z, Westerberg H, Klingberg T (2004). Maturation of white matter is associated with the development of cognitive functions during childhood. *Journal of Cognitive Neuroscience* **16**, 1227–1233.
- Palmen SJ, van Engeland H, Hof PR, Schmitz C (2004). Neuropathological findings in autism. *Brain* **127**, 2572–2583.
- Paus T, Collins DL, Evans AC, Leonard G, Pike B, Zijdenbos A (2001). Maturation of white matter in the human brain: a review of magnetic resonance studies. *Brain Research Bulletin* **54**, 255–266.
- Ramos N, Reichert JG, Smith CJ, Silverman JM, Beshpalova IN, Davis KL, Buxbaum JD (2004). Linkage and association of the mitochondrial aspartate/glutamate carrier SLC25A12 gene with autism. *American Journal of Psychiatry* **161**, 662–669.
- Rose SE, Hatzigeorgiou X, Strudwick MW, Durbridge G, Davies PS, Colditz PB (2008). Altered white matter diffusion anisotropy in normal and preterm infants at term-equivalent age. *Magnetic Resonance Medicine* **60**, 761–767.
- Skranes J, Vangberg TR, Kulseng S, Indredavik MS, Evensen KA, Martinussen M, Dale AM, Haraldseth O, Brubakk AM (2007). Clinical findings and white matter abnormalities seen on diffusion tensor imaging in adolescents with very low birth weight. *Brain* **130**, 654–666.
- Smith SM (2002). Fast robust automated brain extraction. *Human Brain Mapping* **17**, 143–155.
- Villalobos ME, Mizuno A, Dahl BC, Kemmotsu N, Muller RA (2005). Reduced functional connectivity between V1 and inferior frontal cortex associated with visuomotor performance in autism. *Neuroimage* **25**, 916–925.
- Waber DP, De Moor C, Forbes PW, Almli CR, Botteron KN, Leonard G, Milovan D, Paus T, Rumsey J (2007). The NIH MRI study of normal brain development: performance of a population based sample of healthy children aged 6 to 18 years on a neuropsychological battery. *Journal of the International Neuropsychological Society* **13**, 729–746.
- Wassink TH, Hazlett HC, Epping EA, Arndt S, Dager SR, Schellenberg GD, Dawson G, Piven J (2007). Cerebral cortical gray matter overgrowth and functional variation of the serotonin transporter gene in autism. *Archives of General Psychiatry* **64**, 709–717.
- Webb S, Munro CA, Midha R, Stanisiz GJ (2003). Is multicomponent T₂ a good measure of myelin content in peripheral nerve? *Magnetic Resonance Medicine* **49**, 638–645.
- Wechsler D (1999). *Wechsler Abbreviated Scale of Intelligence*. The Psychological Corporation: San Antonio, TX.
- Weng SJ, Wiggins JL, Peltier SJ, Carrasco M, Risi S, Lord C, Monk CS (2010). Alterations of resting state functional connectivity in the default network in adolescents with autism spectrum disorders. *Brain Research* **1313**, 202–214.
- Whittall KP, MacKay AL, Graeb DA, Nugent RA, Li DK, Paty DW (1997). In vivo measurement of T₂ distributions and water contents in normal human brain. *Magnetic Resonance Medicine* **37**, 34–43.
- WHO (2004). *International Statistical Classification of Diseases and Health Related Problems*, 10th revision. World Health Organization: Geneva.
- Wibom R, Lasorsa FM, Tohonen V, Barbaro M, Sterky FH, Kucinski T, Naess K, Jonsson M, Pierri CL, Palmieri F, Wedell A (2009). AGC1 deficiency associated with global cerebral hypomyelination. *New England Journal of Medicine* **361**, 489–495.
- Wolff JJ, Gu H, Gerig G, Elison JT, Styner M, Gouttard S, Botteron KN, Dager SR, Dawson G, Estes AM, Evans AC, Hazlett HC, Kostopoulos P, McKinstry RC, Paterson SJ, Schultz RT, Zwaigenbaum L, Piven J (2012). Differences in white matter fiber tract development present from 6 to 24 months in infants with autism. *American Journal of Psychiatry* **169**, 589–600.
- Woodbury-Smith MR, Robinson J, Wheelwright S, Baron-Cohen S (2005). Screening adults for Asperger Syndrome using the AQ: a preliminary study of its diagnostic validity in clinical practice. *Journal of Autism and Developmental Disorders* **35**, 331–335.