



Parental concerns, socioeconomic status, and the risk of autism spectrum conditions in a population-based study



Xiang Sun^{a,b,c,*}, Carrie Allison^b, Bonnie Auyeung^{b,d}, Simon Baron-Cohen^b, Carol Brayne^a

^a Cambridge Institute of Public Health, Cambridge University School of Medicine, Forvie Site, University of Cambridge, CB2 0SR, UK

^b Autism Research Centre, Department of Psychiatry, University of Cambridge, Douglas House, 18b Trumpington Road, CB2 2AH, UK

^c The Jockey Club School of Public Health and Primary Care, Shatin, NT, Hong Kong

^d Department of Psychology, University of Edinburgh, EH8 9JZ, UK

ARTICLE INFO

Article history:

Received 14 April 2014

Received in revised form 10 July 2014

Accepted 15 July 2014

Available online

Keywords:

Autism Spectrum Conditions

Parental concern

Socioeconomic status

Screening

ABSTRACT

A total number of 11,635 screening packs were distributed to 5–10 year-old children in 136 schools in Cambridgeshire to investigate the associations between levels of parental concern (none/minor/strong), socioeconomic status and the risk of having Autism Spectrum Conditions (ASC). The variables for investigating associations and possible confounders were extracted for analysis, including parental concern question score, SES, age of the child, sex, maternal age at birth, paternal age at birth, mother's age of leaving education, father's age of leaving education, birth order and the number of children in the family. The SES, age of the child, sex and mother's age at leaving education were associated with parental concern. Parents with higher SES reported higher levels of concern (Chi-square = 11.8; $p = 0.02$). However, a higher SES was not associated with the risk of having ASC ($p = 0.50$). After adjusting for potential confounders, the odds of children meeting ASC criteria whose parents had reported strong parental concern were 8.5 times (odds ratio: 8.5; 95%CI: 4.5, 16.2; $p < 0.001$) the odds of children having ASC whose parents reported minor concern. No child met ASC criteria where parents expressed no concerns. Parents with higher social class express more concerns than those from lower social classes. However, the concerns reported by parents in higher SES did not appear to be specific for ASC as there was no relationship between ASC and SES.

© 2014 Elsevier Ltd. All rights reserved.

Introduction

Autism Spectrum Conditions (ASC) refer to the syndrome of childhood onset neuro-developmental disorders characterized by impairments in social interaction and communication, and the presence of repetitive and stereotyped behaviours (Association Psychiatric Association, 2013). In the 1970s, ASC were considered rare, with prevalence estimate of autism around 2 per 10,000 (Gillberg & Wing, 1999; Kawamura, Takahashi, & Ishii, 2008; Waterhouse, 2008). However, the number of children reported as meeting modern ASC criteria has risen dramatically during the last decade (Windham, Zhang,

* Corresponding author at: Cambridge Institute of Public Health, Cambridge University School of Medicine, Forvie Site, University of Cambridge, CB2 0SR, UK. Tel.: +44 01223 763833; fax: +44 01223 330300.

E-mail address: xs227@medschl.cam.ac.uk (X. Sun).

Gunier, Croen, & Grether, 2006). In a UK study, the prevalence of ASC was estimated to be 157 per 10,000 (Baron-Cohen et al., 2009). The most recent Centre of Disease Control (CDC) reported the prevalence of ASC in the US was 1 in 68 (Centre for Disease Control and Prevention, 2014). There are several factors of potential importance which may underlie this increase: (1) changes in diagnostic criteria and screening instruments, (2) changes in study design, (3) more advanced and accessible knowledge about ASC, (4) improvements in public and professional awareness, (5) improved health services and (6) better acceptance of ASC by parents (Baron-Cohen et al., 2009).

Although there has been no definite explanation of the role play between these facts, children with ASC do require more care and support from within and outside families. Parents or those with parental responsibilities play a core role in the life of children with ASC in many regards. During the diagnostic process for ASC, parental recognition plays an important role (Baghdadli, Picot, Pascal, Pry, & Aussilloux, 2003). It can be a crucial factor as early parental awareness is likely to affect the age of recognition of ASC (De & Fombonne, 1998). Reported mean ages of first parental concern range from 14.7 months (Chawarska et al., 2007) to 19.1 months (De & Fombonne, 1998).

By definition, for a diagnosis of ASC, abnormalities should have appeared in early developmental period (Association Psychiatric Association, 2013). However, age at diagnosis is not the same as the age of onset (De & Fombonne, 1998; Howlin & Asgharian, 1999). One study in children with ASC reported that the mean age of first professional advice was 24.1 months. It also reported children with ASC whose parents were concerned about the child's other medical problems had a lower age of first professional advice (De & Fombonne, 1998). The same study found that the age of first parental concern was lower in girls than in boys.

The nature of parental concern has been suggested as potentially useful for clinical case evaluation in child psychiatric referrals (Firth, Grimes, Poppleton, Hall, & Richold, 2000). Studies on the nature of parental concern have indicated that parental concerns about a child's speech and language development are linked to a lower risk of autism than those children whose parents are worried most about other impairments (De & Fombonne, 1998). Two studies found that parents who already had an older child with ASC had been reported to pay more concern about their younger child (between 12 and 36 months) than those with no child with ASC.

Parental concern may also lead to early intervention, which can lead to better outcomes for children with ASC (Fenske, Zalenski, Krantz, & McClannahan, 1985). Children with autism pose considerable behavioural challenges for their parents and other family members on a daily basis due to the nature of their condition. Such children need help to develop skills in establishing their communication, receptive understanding and social interaction behaviours (Blum & Talib, 2006; Rogers, 1998a, 1998b). Chawarska et al. (2007) have suggested that a later age of parental recognition is associated with worse outcomes, although this is yet to be fully confirmed through trials of early intervention. Reviews have indicated that most comprehensive programmes for young children with ASC explicitly involve parents in implementing the strategies (National Research Council, 2001). Studies have reported the potential benefit of parent-implemented early intervention (Rogers et al., 2012). Children in parental training groups have been found to have better social communication skills as rated using the ADI-R (Aldred, Green, & Adams, 2004). Children in parental training groups have been found to score higher on an IQ test than those in intensive applied behaviour analysis (ABA) groups (McConachie & Diggle, 2007).

It has been suggested that improved knowledge and acceptance of autism could be related to the increase in prevalence estimates when there are demands from those children and families (Williams & Brayne, 2006). Such awareness may well be related to socioeconomic status of the parents. Studies based on large population monitoring have provided inconsistent evidence for the association between socioeconomic status and the risk of ASC. In one site of autism and developmental disabilities monitoring (ADDM), using educational attainment as an indicator of socioeconomic status, the prevalence ratio for the highest to lowest education quintile was 2.6:1 (95%CI: 1.6, 4.5) where there was a pre-existing ASC diagnosis (Maenner, Arneson, & Durkin, 2009). In twelve ADDM sites using socioeconomic status (SES), the prevalence ratio for low to medium SES was 0.7 (95%CI: 0.64, 0.76) and that for high to medium SES was 1.25 (95%CI: 1.16, 1.35), regardless of whether a child had a pre-existing ASC diagnosis or not (Durkin et al., 2010). However, in the most recent study in Sweden, children of families with low incomes and parents with manual occupations were found to have a higher risk of ASC (odds ratio = 1.4, 95%CI: 1.3–1.6) (Rai et al., 2012). Sweden has a low inequality index and it is possible that pattern associated with SES might be a better reflection of the population findings rather than an identification of inequality of awareness and access to service.

So far, little research has been conducted to investigate the association between parental concern and the risk of having ASC. The only way to examine the relationship between parental SES, education and the presence of ASC is to conduct the population based study rather than same band analyses or measure parental concern among studies.

Method

Screening instrument

The CAST was developed in the UK, specifically for primary school aged children (aged 4–11), because many children with ASC are often not identified prior to attending primary school (Williams, 2003). The CAST is a 37-item parent completed questionnaire, of which 31 items are scored. One point is assigned for an ASC-positive response and zero for an ASC-negative response on the scored items. Thus, the total score ranges from 0 to 31 (Baron-Cohen et al., 2009). Previous pilot and validation studies have demonstrated the CAST can be used as a screening instrument in large population-based epidemiological research for ASC (Scott, Baron-Cohen, Bolton, & Brayne, 2002b; Williams, Allison et al., 2006; Williams et al.,

2005). Using a cut-off of 15, the sensitivity of the CAST is 100%, specificity is 97%, and positive predictive value (PPV) is 50% (Williams et al., 2005). The psychometric properties of the CAST were investigated in a Chinese population. Two factors were identified including social and communication, and inflexible/stereotyped language and behaviours (Sun et al., 2014).

Questions on parental concern

The SDQ is a relatively brief and user-friendly screening questionnaire for psychosocial problems that can be administered to the parents and teachers of 4 to 17 years old and children themselves aged 11 or over (Goodman, 1997; Goodman, Ford, Simmons, Gatward, & Meltzer, 2000). It has 25 items. The first 20 items measure the following domains: emotional symptoms, conduct problems, hyperactivity and peer problems. The scores are summed to generate a total score ranging from 0 to 40 (Goodman & Goodman, 2012). There have been a number of studies to examine the validity and reliability of the SDQ as a screening instrument for developmental disabilities. A recent review concluded that the internal consistency, test-retest reliability and the inter-rater agreement of the parental version of the SDQ were satisfactory (Stone, Otten, Engels, Vermulst, & Janssens, 2010). The majority of the factor analysis studies of the SDQ found the five-factor solution. They also found the correlations with other measures of psychopathology and the screening ability of the SDQ were sufficient (Stone et al., 2010).

Parental concern was examined by using the other 5 items and 8 sub-questions on the SDQ. Each sub-question has a minimum score of 0 and a maximum score of 3 to categorize the degree of parental concern about their child (Table 1). The maximum score for parental concern is 24.

The first question asks whether parents had concerns about their child in terms of their emotions, concentration, behaviour or getting along with others. If the parent answers no (score 0: indicating the parent had no concerns about their child), they did not need to continue through the whole questionnaire. The scores for the parental concern questions are divided into three categories: no concern = 0, minor concern = 1–12, great concern = 13–24.

Study design and sample

In this study, data on parental concern and the risk of ASC were collected during a prospective screening study, the Social and Communication Research and Epidemiology study (SCORE). The study had full ethical approval from the Ethics Committee of the University. The SCORE study used the Childhood Autism Spectrum Test (CAST) to screen for ASC in 5–10 year-old children in Cambridgeshire. In total, 136 mainstream schools within the county of Cambridgeshire, including Cambridge City, East Cambridgeshire, South Cambridgeshire and Fenland, were invited to participate in this study (Baron-Cohen et al., 2009). The first section of the CAST pack consisted of the CAST (37 items), some questions asking about medical conditions or developmental disorders (15 items) and parental concern questions from the SDQ (Goodman, 1997). Section 2 consisted of questions about the parents. Section 3 consisted of questions about the education and occupations of the

Table 1
Questions on parental concern in the screening pack.

No	Question wording	Score = 0 (N %)	Score = 1 (N %)	Score = 2 (N %)	Score = 3 (N %)	Missing (N %)	NA (N %)
1	Overall, do you think your child has difficulties in one or more of the following areas: emotions, concentration, behaviour or being able to get on with other people?	2262 (66.5%)	814 (23.9%)	239 (7.0%)	37 (1.1%)	52 (1.5%)	0
2	How long have these difficulties been present?	14 (0.4%)	57 (1.7%)	115 (3.4%)	903 (26.5%)	53 (1.6%)	2262 (66.5%)
3	Do the difficulties upset or distress your child?	293 (8.6%)	514 (15.1%)	226 (6.6%)	54 (1.6%)	54 (1.6%)	2263 (66.5%)
4	Do the difficulties interfere with your child's everyday life in home life?	280 (8.2%)	540 (15.9%)	210 (6.2%)	62 (1.8%)	49 (1.4%)	2263 (66.5%)
5	Do the difficulties interfere with your child's everyday life in friendship?	467 (13.7%)	418 (12.3%)	154 (4.5%)	50 (1.5%)	52 (1.5%)	2263 (66.5%)
6	Do the difficulties interfere with your child's everyday life in classroom learning?	284 (8.3%)	466 (13.7%)	239 (7.0%)	88 (2.6%)	64 (1.9%)	2263 (66.5%)
7	Do the difficulties interfere with your child's everyday life in leisure activities?	539 (15.9%)	387 (11.4%)	129 (3.8%)	30 (0.9%)	56 (1.7%)	2263 (66.5%)
8	Do the difficulties put a burden on you or the family as a whole?	339 (10.0%)	504 (14.8%)	176 (5.2%)	70 (2.0%)	54 (1.6%)	2261 (66.4%)

Q1: 0 = No; 1 = minor difficulties; 2 = definite difficulties; 3 = severe difficulties; Q2: 0 = Less than a month; 1 = less than 5 months; 2 = 6–12 months; 3 = over a year; Q3–8: 0 = not at all; 1 = only a little; 2 = quite a lot; 3 = a great deal; NA = not applicable.

parents which were used to categorize their socioeconomic status. After the collection of the CAST pack, the children were divided into three score groups (≤ 11 , 12–14, ≥ 15). All the children above the cut-point of 15 (≥ 15) and a randomly selected 33% of the borderline group (12–14) were invited for a detailed diagnostic assessment to identify potential cases of ASC in the general population. No child in low score group (≤ 11) was invited because no children with ASC had been identified among low score group, no children in low score group were diagnosed with ASC (Scott, Baron-Cohen, Bolton, & Brayne, 2002a; Williams et al., 2005). The diagnostic battery of the detailed assessment included the Autism Diagnostic Observational Schedule (ADOS; Lord, Rutter, DiLavore, & Risi, 2001) (ADOS) and the Autism Diagnostic Interview-Revised (ADI-R; Lord, Rutter, & Le Couteur, 1994).

A total of 11,635 CAST packs were distributed via class registration and returned to the research centre by parents. The pack was distributed to participating schools in six batches between February 2003 and March 2004. In total, 3404 questionnaires (response rate = 29%) were returned for analysis.

Confirmation of children with ASC

Cases were identified through two different routes: the questionnaires through mainstream schools and records from Special Educational Needs (SEN) registration. Population-based screening was conducted in Cambridgeshire. Children who were reported having a diagnosis of ASC based on their records in the SEN register received an estimation of their ASC diagnostic sub-grouping using ICD-10 (World Health Organisation, 1993). Those children were considered to be cases without further assessment.

After screening, the ADOS and ADI-R were used as diagnostic instruments. Previous research suggested that using the ADI-R and ADOS alone can be less reliable in detecting milder cases on the autism spectrum. Because the agreement between the ADOS and ADI-R is not 100% and the ADI-R only provides diagnostic cut-off for classic autism (de et al., 2004; Couteur, Haden, Hammal, & McConachie, 2008). Thus, a consensus diagnosis based on information gathered through the ADOS and ADI-R and the clinical judgement of clinical members on the research team was adopted. The following subtypes were included into the diagnosis of ASC: classic autism, Asperger syndrome and PDD-NOS. For all children who were given a diagnosis of ASC, their assessment data was re-examined and reviewed according to ICD-10 (World Health Organisation, 1993). The ADOS and ADI-R were administered by members of the research team who were fully trained in the use of these instruments to reliability levels for research purposes (Baron-Cohen et al., 2009). The examiners had been working in our research centre for a number of validation and prevalence studies (Allison et al., 2007; Scott et al., 2002a, 2002b; Williams et al., 2005; Williams, Higgins, & Brayne, 2006).

Statistical analyses

Analyses were conducted using STATA 10.0 statistical software. In order to examine the potential confounding effects, the following variables were extracted from the dataset: an anonymous identification number for each child, age of the child, sex, parental concern question score, socioeconomic status, maternal age at birth, paternal age at birth, mother's age of leaving education father's age of leaving education, birth order and the number of children in the family.

The socioeconomic status of this sample was generated using the self-coded National Statistics Socio-economic Classification (NS-SEC) as follows (Office for National Statistics): (1) Managerial and professional occupations, (2) Intermediate occupations, (3) Small employers and own account workers, (4) Lower supervisory and technical occupations and (5) Semi-routine and routine occupations. The lowest three categories were combined, thus there were three categories: Manager, Intermediate and Lower.

The distributions of all the variables were examined using histograms or box plots and the following were categorized: parental concern, maternal age at birth, paternal age at birth, mother's age of leaving education, father's age of leaving education.

The associations between parental concern and each of the other variables were examined using chi-square tests. The unadjusted associations between each variable and the risk of ASC were examined using logistic regression. The variables that associated with both ASC and parental concern were put into an unconditional logistic regression model. The association between parental concern and the risk of ASC was examined by adjusting all the other variables in the model using unconditional logistic regression. The interaction between parental and other variables was examined. The odds ratios and 95% confidence intervals (CI) and *p*-values were computed for all variables.

Results

Characteristics of participants

Within the 3404 returned questionnaires, 3329 records (97.8%) were available for analysis and 75 records (2.2%) were excluded due to missing data on the SDQ. After screening and standardized diagnostic assessment in mainstream schools, 9 children from mainstream schools were given a research diagnosis of ASC. There were 37 children who were reported to have an existing diagnosis of ASC according to schools' SEN registers. Thus, altogether there were 46 children with ASC and 3283 children without ASC.

A total of 2260 parents (66.4%) indicated that they had no concerns about their child, 866 (25.4%) had minor concerns and 226 (6.6%) had great concerns. In 52 (1.5%) questionnaires, the data on parental concern were missing. The process of case identification is shown in Fig. 1. The characteristics of the children with ASC and without ASC are shown in Table 2.

Parental concern and other variables

The Chi-square test indicated four variables that were associated with parental concern in this sample (Table 3): age, sex, socioeconomic status and mother's age of leaving education.

Association between all variables and the risk of ASC

Compare children with ASC with children without ASC (the unadjusted odds ratio), three out of ten variables were associated with the risk of ASC: sex, parental concern and paternal age at birth. The distribution of sex between cases and




Fig. 1. Flow chart of case identification.

Table 2
Characteristics of cases and controls in the main analysis.

Variable	Category	Cases (N = 46)	(%)	Controls (N = 3283)	(%)
Age group (yrs)	4–5	8	(17.4)	341	(10.4)
	6	8	(17.4)	654	(20.0)
	7	12	(26.1)	866	(26.4)
	8	8	(17.4)	827	(25.2)
	9–10	10	(21.7)	571	(17.4)
Sex	Male	35	(76.1)	1623	(49.4)
	Female	11	(23.9)	1652	(50.3)
Parental concern	No concern	0	(0.0)	2237	(68.6)
	Minor concern	15	(33.3)	468	(25.6)
	Great concern	30	(66.7)	189	(5.8)
	Missing	1	(2.2)	21	(0.6)
Socioeconomic status	Manager	17	(37.0)	1479	(45.1)
	Intermediate	10	(21.7)	548	(16.7)
	Lower	10	(21.7)	677	(20.6)
	Missing	9	(19.6)	579	(17.6)
Maternal age	≤29	20	(1.3)	1462	(97.3)
	30–34	15	(1.3)	1161	(97.2)
	≥35	11	(1.8)	613	(97.5)
Paternal age	≤29	15	(1.8)	788	(96.7)
	30–34	7	(0.6)	1089	(98.0)
	≥35	20	(2.1)	955	(96.6)
Mother's age of leaving education	≤18	31	(1.6)	1943	(97.0)
	19–22	7	(0.8)	813	(97.8)
	≥23	5	(1.4)	354	(98.1)
Father's age of leaving education	≤18	30	(1.8)	1661	(96.8)
	19–22	4	(0.6)	646	(98.5)
	≥23	5	(1.2)	422	(97.5)
Birth order	1	22	(1.4)	1518	(96.8)
	2	16	(1.4)	1120	(96.7)
	≥3	8	(1.3)	586	(96.4)
No. of children	1–2	28	(60.9)	1977	(60.2)
	≥3	18	(39.1)	1285	(39.1)
	Missing	0	(0)	21	(0.6)

Parental concern: No concern: Score 0; Minor concern: Score 1–12; Great concern: 12–24.

controls was different. The sex ratio (boys Vs girls) was more than 3:1, which showed a predominance of ASC in boys. The crude odds ratio of having autism among boys and girls was 3.2 (95%CI = 1.6, 6.4; $p = 0.001$). There was a highly significant association between gender and the risk of having ASC. As there were no cases in the first category of parental concern (the no concerns group), the second category (minor concerns) was used as the reference to investigate the association between the degree of parental concern and the risk of ASC. The odds of having ASC among children whose parents had great concerns (scored 13–24) were 8.8 times greater than for children whose parents had minor concerns (scored >12) (crude odds ratio = 8.8; 95% CI: 4.7, 16.8; $p < 0.001$). This suggests a highly significant association between parental concern and the risk of ASC. The higher the parental concern score, the more likely that the child would meet ASC criteria. The odds of having ASC when the paternal age was between 30 and 34 at the time of the child's birth were significantly lower than for those whose fathers were younger than 30 when the child was born (crude odds ratio = 0.3; 95%CI = 0.1, 0.9). The results of unadjusted association are shown in [Table 4](#).

Adjusted model

Only variables that were significantly associated with both the risk of ASC and parental concern were included in the unconditional logistic regression model. Thus, there were two variables in the model: sex and parental concern ([Table 5](#)). After adjustment for sex, the odds ratio for parental concern reduced to 8.5 (95%CI: 4.5, 16.2; $p < 0.001$), so hardly changed (8.8–8.5).

For boys, the odds of having ASC among children whose parents had great concerns were 27.6 times greater than for children whose parents had minor concerns (95%CI: 5.8, 131.2; $p < 0.001$). In girls, however, the odds ratio between children whose parents had great concerns and those whose parents had minor concerns was lower at 6.0 (95%CI: 2.9, 12.4; $p < 0.001$). The odds of being a boy with autism were 1.7 times greater than the odds of being a girl with autism (95%CI: 0.8, 3.4; $p = 0.15$). The interaction between parental concern and sex was found to fall short of significance ($p = 0.08$).

Table 3
Association between variables and parental concern.

Variable	Parental concern	
	Chi-square	P-value
Age	16.0	0.04
Sex	87.5	<0.001
Socioeconomic status	11.8	0.02
Maternal age at birth	2.8	0.59
Paternal age at birth	8.1	0.09
Mother's age of leaving education	15.0	0.005
Father's age of leaving education	7.8	0.10
Birth order	7.0	0.13
Number of children	0.3	0.85

In this final model, only parental concern was significantly associated with the risk of ASC. Thus, there was a highly statistically significant association between parental concern and the risk of ASC. This association was independent of socioeconomic status of parents and other potential confounders.

Discussion

In this study, the degree of parental concern in a large population sample was determined through questions on the SDQ. A higher score on these questions indicated that parents considered their children's difficulties had more severe impact on their daily lives. The aim of this study was to explore the association between parental concern and the risk of having ASC. It was therefore possible to investigate whether more parental concern about their child's emotions, behaviour and getting on

Table 4
Unadjusted association between variables and the risk of ASC in the main analysis.

Variable	Category	UOR	95% CI	P-value
Age group (yrs)	4–5	1.0	Reference	0.45
	6	0.5	(0.2, 1.4)	
	7	0.6	(0.2, 1.5)	
	8	0.4	(0.2, 1.1)	
	9–10	0.7	(0.3, 1.9)	
Sex	Female	1.0	Reference	0.001
	Male	3.2	(1.6, 6.4)	
Parental concern	No concern	–	–	–
	Minor concern	1.0	Reference	<0.001
	Great concern	8.8	(4.7, 16.8)	
Socioeconomic status	Manager	1.0	Reference	0.50
	Intermediate	1.6	(0.7, 3.5)	
	Lower	1.3	(0.6, 2.8)	
Maternal age at birth	≤29	1.0	Reference	0.69
	30–34	0.9	(0.5, 1.9)	
	≥35	1.3	(0.6, 2.8)	
Paternal age at birth	≤29	1.0	Reference	0.02
	30–34	0.3	(0.1, 0.8)	
	≥35	1.1	(0.6, 2.2)	
Mother's age of leaving education	≤18	1.0	Reference	0.34
	19–22	0.5	(0.2, 1.2)	
	≥23	0.9	(0.3, 2.3)	
Father's age of leaving education	≤18	1.0	Reference	0.11
	19–22	0.3	(0.1, 0.9)	
	≥23	0.7	(0.3, 1.7)	
Birth order	1	1.0	Reference	0.99
	2	1.0	(0.5, 1.9)	
	≥3	0.9	(0.4, 2.1)	
No. of children	1–2	1.0	Reference	0.97
	≥3	1.0	(0.5, 1.8)	

Parental concern: No concern: Score 0; Minor concern: Score 1–12; Great concern: 13–24
UOR: unadjusted odds ratio.

Table 5
Adjusted model in the main analysis.

Variable	AOR	95%CI	P-value
Sex			
Female	1.0	Reference	0.15
Male	1.7	(0.8, 3.4)	
Parental concern			
No concern	–	–	
Minor concern	1.0	Reference	<0.001
Great concern	8.5	(4.5, 16.2)	

Parental concern: No concern: Score 0; Minor concern: Score 1–12; Great concern: 13–24.
AOR: Adjusted for all variables in the table.

with others would lead to a higher chance of the child meeting ASC criteria for a diagnosis. This was the case in this sample. Socioeconomic status was also associated with parental concerns but not related to ASC diagnosis.

Limitations

There are a number of limitations of this study. The response rate for the screening study was low (29%), similar to other parent-report screening studies (Baron-Cohen et al., 2009; Williams, Allison et al., 2006). Since autism is a relatively rare condition, a few missing cases in non-responders could influence the results. In this study, differential response according to SES, parental concerns and ASC presence could impact the association.

There may be recall bias among parents. Parents answer the questions based on their memories, and parents whose children meeting ASC even if not diagnosed, might have had a much clearer memory for certain periods during their child's early development than those parents whose children do not meet the condition's criteria. The questions themselves could be seen to be part of the same interview and thus it is to be expected that there is a close relationship.

Since not all children who scored 12–14 on the CAST were invited to further assessment there may be potential cases among these children. In this study these potential children with ASC were considered as non-ASC. This could have resulted in an underestimation of the number of children with ASC and overestimation of the number of children without ASC. However, only one child who attended further assessment after scoring between 12 and 14 met criteria for ASC diagnosis. Thus, it is unlikely this could have greatly influenced the results.

In the diagnostic phase of this study, cognitive measurements or IQ tests were not adopted other than the two diagnostic instruments. Such measurements would have helped to better understand and recognize the children, especially those who were given a diagnosis of ASC. However, as these instruments are not for diagnosis, it is not likely their results could substantially influence the final consensus diagnosis.

There were only 46 cases compared to 3283 controls during the unconditional logistic analysis and some reference groups (such as parental concern) had very small numbers. For a case–control study, the numbers should be equal to or larger than five in each group, so this might have partly exaggerated the effect of the risk factor. However, the purpose of this paper is not to conduct a case–control study which would require an even or similar children with ASC and children without ASC. The purpose was to intend to understand what would be the association between parental concerns, socioeconomic status, and the risk of having ASC in a general population. An even sample design would not provide sufficient or reliable information for our question. It is admitted that there is a large difference in the number of children with ASC and children without ASC in our sample. However, since the generally agreed prevalence of ASC in the general population is around 1% (Centre for Disease Control and Prevention, 2012; Fombonne, 2009), the prevalence of our sample should be acceptable and reasonable to fulfil the purpose of study design (prevalence of this study is around 1.4%). As the basic question is to understand the situation in a general population, the whole population sample was adopted. The statistical analyses were conducted within this sample rather than picking up fewer children without ASC to compare with children with ASC. Thus, to speak from a general population public health of view, using the sample from a general population would be more sensible than only comparing even cases and controls. Since the number of cases depends on the prevalence of ASC, to have more children with ASC in such study, much larger sample sizes are needed in the future.

Cambridgeshire might not be representative of the UK as a whole, so caution should be employed when considering the national generalisability of these results. For example, SES and parental education in Cambridgeshire are generally higher than in other regions of the UK (Baron-Cohen et al., 2009).

Parental concern and public health implications

According to these data, there were no cases in children whose parents who did not report concerns. In children whose parents who did report concerns, there was a highly significant association between parent-reported concern and the risk of ASC. As the closest carers of the child, parents have the most opportunities to recognize abnormalities in their children, but may themselves not easily pick up behaviours and styles which others may consider diagnostic.

Many clinical features of ASC gradually come to the notice of parents before the second birthday of the child (Chawarska et al., 2007). Children with ASC usually experience direct and indirect consequences related to social interaction and communication deficits. These social impairments are diverse and can involve speech, linguistic conventions and interpersonal interaction. Childhood serves to develop these skills in order to progress throughout the life course. If autistic children could be identified earlier and given proper help and guidance in their development, their outcomes might be improved. With this thought in mind, researchers have done a lot to achieve this goal. Both randomized controlled trials and controlled studies have been carried out in this field (Aldred, Green, Emsley, & McConachie, 2012; Howlin, Magiati, & Charman, 2009; Virues-Ortega, 2010). Outcomes of interventions have been measured with direct testing of socio-communication skills and IQ, as well as parents' and teacher's reports of adaptive skills and behavioural problems (McConachie & Diggle, 2007). This highly significant association between parental concern and autism risk, as well as the evidence on the effectiveness of parent-participated intervention and surveillance, suggest that this is an area for further exploration.

Parental concern and screening for ASC

Screening is the prospective identification of unrecognized disorders through the application of specific tests or examinations. Screening for ASC has been recommended in the USA for early detection with the goal of obtaining improved outcomes (Levy, Mandell, & Schultz, 2009). There is increasing agreement amongst clinicians that intervention in ASC is more effective if it commences before the age of four (Baron-Cohen et al., 2000; Dumont-Mathieu & Fein, 2005; Tanguay, 2000). Previous studies have provided evidence that combining screening instruments with asking about parental concerns can improve the utility and efficiency of the instruments (Baird et al., 2001; Glascoe, 1997; Glascoe, 2003; Glascoe & Sandler, 1995). Dietz's study on the screening programme for ASC in the Netherlands found that a large number of children were "lost" at each stage of the screening procedure because parents did not want to cooperate (Dietz, Swinkels, van Daalen, van Engeland, & Buitelaar, 2006). It also suggested that hesitation among parents could be reflected in the delay between screening and further examination of children. Sugiyama's study in Japan found that toddlers who failed a developmental check up at 18 months, and whose parents refused further investigations, showed more developmental problems at the 3-year follow up than the children who did participate in further investigations (Sugiyama & Abe, 1989). The present study included questions on parental concerns in screening in a general population in the UK. Our findings suggest that combining the observations of parents with the employment of screening tests might be valuable for early detection of potential cases of ASC.

Sex and parental concern

Sex was associated with both parental concern and the risk of ASC. The higher prevalence of ASC in boys compared to girls has been reported by previous studies and the ratio is around 4:1 (Fombonne, 2009; Volkmar, Szatmari, & Sparrow, 1993). In this study, the odds of a boy having a diagnosis of ASC were 27.6 times higher in those children whose parental concern was great compared with those parents had minor concerns.

After adjustment for sex, the association between parental concern and the risk of ASC hardly changed.

Socioeconomic status and parental concern

Research regarding a possible association between socioeconomic status or social class and the risk of ASC can be traced back to 1943, when Kanner first described autism. He characterized high-risk families as highly intelligent and educated with good employment. Since then, many studies have used parental occupation, educational background or the level of intelligence to generate social class categories (Croen, Grether, & Selvin, 2002).

Early studies have agreed with Kanner's description (Hoshino, Kumashiro, Yashima, Tachibana, & Watanabe, 1982), while others have found no such association (Cialdella & Mamelle, 1989; Croen et al., 2002; Cryan, Byrne, O'Donovan, & O'Callaghan, 1996; Hoshino et al., 1982), suggesting that higher social class is not related to the risk of autism. Recent studies on the association between socioeconomic status and the risk of autism reported conflicted results (Durkin et al., 2010; King & Bearman, 2011; Rai et al., 2012). Inequality in access to service due to different SES status of parents is not unusual. Some researchers have argued that this observed association could also be explained by other factors such as better diagnosis and early referral to health institutions because parents with a higher socioeconomic status may have more access to health services (Croen et al., 2002; Wing, 1980). Other studies have suggested that the association results from different perceptions of autism among the highly educated professionals and parents (Bhasin & Schendel, 2007; Cuccaro et al., 1996). The present study found that socioeconomic status was associated with parental concern but not with the risk of ASC in this population. If socioeconomic status is a confounder, it should be associated with both the risk of ASC and parental concern. Thus, the hypothesis that socioeconomic status might confound the association between parental concern and the risk of ASC is not supported by this study.

Conclusion

Parents with higher social class express more concerns than those from lower social classes. However, the concerns reported by parents in higher SES did not appear to be specific for ASC as there was no relationship between ASC and SES.

Parental concern itself was strongly associated with a child meeting ASC criteria. The higher the degree of parental concern over their child in terms of emotions, concentration, behaviour or getting on with others, the more likely that the child will have ASC, independently of potential confounders. Conversely, this study shows that where there is no parental concern expressed a child is extremely unlikely to meet diagnostic criteria. These findings should be of value in discussions related to which measures at what times are helpful in identifying children with ASC.

Acknowledgements

We are grateful to all families for their participation in this study. We are indebted to Stephen Sharp and Yu-tzu Wu for the discussions on data analysis. XS was supported by the Waterloo Foundation, Cambridge Commonwealth Trust, Clare Hall of the University of Cambridge and Great Britain-China Educational Trust during initial writing up and then funded by the International Development Fund Cambridge-CUHK Collaboration on Autism Research in Hong Kong and China during the later writing up of this paper. CA, SBC, and BA were supported by the MRC UK. This study was conducted in association with the NIHR CLAHRC for Cambridgeshire and Peterborough NHS Foundation Trust.

References

- Aldred, C., Green, J., & Adams, C. (2004). A new social communication intervention for children with autism: Pilot randomised controlled treatment study suggesting effectiveness. *Journal of Child Psychology and Psychiatry*, 45(8), 1420–1430.
- Aldred, C., Green, J., Emsley, R., & McConachie, H. (2012). Brief report: Mediation of treatment effect in a communication intervention for pre-school children with autism. *Journal of Autism and Developmental Disorders*, 42(3), 447–454.
- Allison, C., Williams, J., Scott, F., Stott, C., Bolton, P., & Baron-Cohen, S. (2007). The Childhood Asperger Syndrome Test (CAST): Test–retest reliability in a high scoring sample. *Autism*, 11(2), 177–190.
- Association Psychiatric Association (2013). *Diagnostic and statistical manual of mental disorders (DSM-5)* (5th ed.). Washington, DC: American Psychiatric Association.
- Baghdadli, A., Picot, M. C., Pascal, C., Pry, R., & Aussilloux, C. (2003). Relationship between age of recognition of first disturbances and severity in young children with autism. *European Child and Adolescent Psychiatry*, 12(3), 122–127.
- Baird, G., Charman, T., Cox, A., Baron-Cohen, S., Swettenham, J., & Wheelwright, S. (2001). Current topic: Screening and surveillance for autism and pervasive developmental disorders. *Archives of Diseases in Childhood*, 84(6), 468–475.
- Baron-Cohen, S., Scott, F. J., Allison, C., Williams, J., Bolton, P., & Matthews, F. E. (2009). Prevalence of autism-spectrum conditions: UK school-based population study. *British Journal of Psychiatry*, 194(6), 500–509.
- Baron-Cohen, S., Wheelwright, S., Cox, A., Baird, G., Charman, T., & Swettenham, J. (2000). Early identification of autism by the Checklist for Autism in Toddlers (CHAT). *Journal of Royal Society of Medicine*, 93(10), 521–525.
- Bhasin, T. K., & Schendel, D. (2007). Sociodemographic risk factors for autism in a US metropolitan area. *Journal of Autism and Developmental Disorders*, 37(4), 667–677.
- Blum, J. D., & Talib, N. (2006). Balancing individual rights versus collective good in public health enforcement. *Medicine and Law*, 25(2), 273–281.
- Centre for Disease Control and Prevention (2012). Prevalence of autism spectrum disorders – Autism and Developmental Disabilities Monitoring Network, 14 sites, United States, 2008. *MMWR Surveillance Summaries*, 61(3), 1–19.
- Centre for Disease Control and Prevention (2014). Prevalence of autism spectrum disorder among children aged 8 years – Autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *MMWR Surveillance Summaries*, 63(Suppl 2), 1–21.
- Chawarska, K., Paul, R., Klin, A., Hannigen, S., Dichtel, L. E., & Volkmar, F. (2007). Parental recognition of developmental problems in toddlers with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 37(1), 62–72.
- Cialdella, P., & Mabelle, N. (1989). An epidemiological study of infantile autism in a French department (Rhône): A research note. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 30(1), 165–175.
- Croen, L. A., Grether, J. K., & Selvin, S. (2002). Descriptive epidemiology of autism in a California population: Who is at risk? *Journal of Autism and Developmental Disorders*, 32(3), 217–224.
- Cryan, E., Byrne, M., O'Donovan, A., & O'Callaghan, E. (1996). A case-control study of obstetric complications and later autistic disorder. *Journal of Autism and Developmental Disorders*, 26(4), 453–460.
- Cuccaro, M. L., Wright, H. H., Rownd, C. V., Abramson, R. K., Waller, J., & Fender, D. (1996). Professional perceptions of children with developmental difficulties: The influence of race and socioeconomic status. *Journal of Autism and Developmental Disorders*, 26(4), 461–469.
- de, B. A., Sytema, S., Ketelaars, C., Kraijer, D., Mulder, E., & Volkmar, F. (2004). Interrelationship between Autism Diagnostic Observation Schedule-Generic (ADOS-G), Autism Diagnostic Interview-Revised (ADI-R), and the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) classification in children and adolescents with mental retardation. *Journal of Autism and Developmental Disorders*, 34(2), 129–137.
- De, G., & Fombonne, E. (1998). Parental recognition of developmental abnormalities in autism. *European Child & Adolescent Psychiatry*, 7(3), 131–136.
- Dietz, C., Swinkels, S., van Daalen, E., van Engeland, H., & Buitelaar, J. K. (2006). Screening for autistic spectrum disorder in children aged 14–15 months. II: Population screening with the Early Screening of Autistic Traits Questionnaire (ESAT). Design and general findings. *Journal of Autism and Developmental Disorders*, 36(6), 713–722.
- Dumont-Mathieu, T., & Fein, D. (2005). Screening for autism in young children: The Modified Checklist for Autism in Toddlers (M-CHAT) and other measures. *Mental Retardation and Developmental Disabilities Research Reviews*, 11(3), 253–262.
- Durkin, M. S., Maenner, M. J., Meaney, F. J., Levy, S. E., DiGuseppi, C., & Nicholas, J. S. (2010). Socioeconomic inequality in the prevalence of autism spectrum disorder: Evidence from a U.S. cross-sectional study. *PLOS ONE*, 5(7), e11551.
- Fenske, E. C., Zalski, S., Krantz, P. J., & McClannahan, L. E. (1985). Age at intervention and treatment outcome for autistic-children in a comprehensive intervention program. *Analysis and Intervention in Developmental Disabilities*, 5(1–2), 49–58.
- Firth, H., Grimes, A., Poppleton, H., Hall, R., & Richold, P. (2000). Assessment of parents' concerns and evaluation of outcomes. *Journal of Public Health Medicine*, 22(4), 473–478.
- Fombonne, E. (2009). Epidemiology of pervasive developmental disorders. *Pediatrics in Review*, 65(6), 591–598.
- Gillberg, C., & Wing, L. (1999). Autism: Not an extremely rare disorder. *Acta Psychiatrica Scandinavica*, 99(6), 399–406.
- Glascoc, F. P. (1997). Parents' concerns about children's development: Prescreening technique or screening test? *Pediatrics*, 99(4), 522–528.
- Glascoc, F. P. (2003). Parents' evaluation of developmental status: How well do parents' concerns identify children with behavioral and emotional problems? *Clinical Pediatrics (Phila)*, 42(2), 133–138.
- Glascoc, F. P., & Sandler, H. (1995). Value of parents' estimates of children's developmental ages. *Journal of Pediatrics*, 127(5), 831–835.
- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38(5), 581–586.
- Goodman, A., & Goodman, R. (2012). Strengths and Difficulties Questionnaire scores and mental health in looked after children. *British Journal of Psychiatry*, 200(5), 426–427.

- Goodman, R., Ford, T., Simmons, H., Gatward, R., & Meltzer, H. (2000). Using the Strengths and Difficulties Questionnaire (SDQ) to screen for child psychiatric disorders in a community sample. *British Journal of Psychiatry*, 177, 534–539.
- Hoshino, Y., Kumashiro, H., Yashima, Y., Tachibana, R., & Watanabe, M. (1982). The epidemiological study of autism in Fukushima-ken. *Folia Psychiatrica et Neurologica Japonica*, 36(2), 115–124.
- Howlin, P., & Asgharian, A. (1999). The diagnosis of autism and Asperger syndrome: Findings from a survey of 770 families. *Developmental Medicine and Child Neurology*, 41(12), 834–839.
- Howlin, P., Magiati, I., & Charman, T. (2009). Systematic review of early intensive behavioral interventions for children with autism. *Journal of Intellectual & Developmental Disability*, 114(1), 23–41.
- Kawamura, Y., Takahashi, O., & Ishii, T. (2008). Reevaluating the incidence of pervasive developmental disorders: Impact of elevated rates of detection through implementation of an integrated system of screening in Toyota, Japan. *Journal of Neuropsychiatry & Clinical Neuroscience*, 62(2), 152–159.
- King, M. D., & Bearman, P. S. (2011). Socioeconomic status and the increased prevalence of autism in California. *American Sociological Review*, 76(2), 320–346.
- Le Couteur, A., Haden, G., Hammal, D., & McConachie, H. (2008). Diagnosing autism spectrum disorders in pre-school children using two standardised assessment instruments: The ADI-R and the ADOS. *Journal of Autism and Developmental Disorders*, 38(2), 362–372.
- Levy, S. E., Mandell, D. S., & Schultz, R. T. (2009). Autism. *Lancet*, 374(9701), 1627–1638.
- Lord, C., Rutter, M., DiLavore, P., & Risi, S. (2001). *Autism diagnostic observation schedule (ADOS)*. Los Angeles, CA: Western Psychological Services.
- Lord, C., Rutter, M., & Le Couteur, A. (1994). Autism Diagnostic Interview-Revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism Developmental Disorders*, 24(5), 659–685.
- Maenner, M. J., Arneson, C. L., & Durkin, M. S. (2009). Socioeconomic disparity in the prevalence of autism spectrum disorder in Wisconsin. *World Medical Journal*, 108(5), 253–255.
- McConachie, H., & Diggle, T. (2007). Parent implemented early intervention for young children with autism spectrum disorder: A systematic review. *Journal of Evaluation in Clinical Practice*, 13(1), 120–129.
- National Research Council (2001). *Educating children with autism. Committee on educational intervention for children with autism. Commission on behavioural and social science and education*. National Academy Press.
- Office for National Statistics (2002). *National statistics Socio-economic Classification (NS-SEC)*.
- Rai, D., Lewis, G., Lundberg, M., Araya, R., Svensson, A., & Dalman, C. (2012). Parental socioeconomic status and risk of offspring autism spectrum disorders in a Swedish population-based study. *Journal of the American Academy of Child & Adolescent Psychiatry*, 51(5), 467–476.
- Rogers, S. J. (1998a). Empirically supported comprehensive treatments for young children with autism. *Journal of Clinical Child Psychology*, 27(2), 168–179.
- Rogers, S. J. (1998b). Neuropsychology of autism in young children and its implications for early intervention. *Mental Retardation and Developmental Disabilities Research Reviews*, 4(2), 104–112.
- Rogers, S. J., Estes, A., Lord, C., Vismara, L., Winter, J., & Fitzpatrick, A. (2012). Effects of a brief Early Start Denver model (ESDM)-based parent intervention on toddlers at risk for autism spectrum disorders: A randomized controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry*, 51(10), 1052–1065.
- Scott, F. J., Baron-Cohen, S., Bolton, P., & Brayne, C. (2002a). Brief report: Prevalence of autism spectrum conditions in children aged 5–11 years in Cambridgeshire, UK. *Autism*, 6(3), 231–237.
- Scott, F. J., Baron-Cohen, S., Bolton, P., & Brayne, C. (2002b). The CAST (childhood Asperger syndrome test): Preliminary development of a UK screen for mainstream primary-school-age children. *Autism*, 6(1), 9–31.
- Stone, L. L., Otten, R., Engels, R. C., Vermulst, A. A., & Janssens, J. M. (2010). Psychometric properties of the parent and teacher versions of the strengths and difficulties questionnaire for 4- to 12-year-olds: A review. *Clinical Child and Family Psychology Review*, 13(3), 254–274.
- Sugiyama, T., & Abe, T. (1989). The prevalence of autism in Nagoya, Japan: A total population study. *Journal of Autism and Developmental Disorders*, 19(1), 87–96.
- Sun, X., Allison, C., Auyeung, B., Matthews, F. E., Norton, S., & Baron-Cohen, S. (2014). Psychometric properties of the mandarin version of the childhood autism spectrum test (CAST): An exploratory study. *Journal of Autism and Developmental Disorders*, 44(7), 1565–1576.
- Tanguay, P. E. (2000). Pervasive developmental disorders: A 10-year review. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39(9), 1079–1095.
- Virues-Ortega, J. (2010). Applied behavior analytic intervention for autism in early childhood: Meta-analysis, meta-regression and dose-response meta-analysis of multiple outcomes. *Clinical Psychology Review*, 30(4), 387–399.
- Volkmar, F. R., Szatmari, P., & Sparrow, S. S. (1993). Sex differences in pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 23(4), 579–591.
- Waterhouse, L. (2008). Autism overflows: Increasing prevalence and proliferating theories. *Neuropsychology Review*, 18(4), 273–286.
- Williams, J. (2003). *Screening for autism spectrum disorders*. University of Cambridge.
- Williams, J., Allison, C., Scott, F., Stott, C., Bolton, P., & Baron-Cohen, S. (2006a). The childhood Asperger syndrome test (CAST): Test-retest reliability. *Autism*, 10(4), 415–427.
- Williams, J., & Brayne, C. (2006). Screening for autism spectrum disorders: What is the evidence? *Autism*, 10(1), 11–35.
- Williams, J. G., Higgins, J. P., & Brayne, C. E. (2006). Systematic review of prevalence studies of autism spectrum disorders. *Archives of Diseases in Childhood*, 91(1), 8–15.
- Williams, J., Scott, F., Stott, C., Allison, C., Bolton, P., & Baron-Cohen, S. (2005). The CAST (childhood Asperger syndrome test): Test accuracy. *Autism*, 9(1), 45–68.
- Windham, G. C., Zhang, L., Gunier, R., Croen, L. A., & Grether, J. K. (2006). Autism spectrum disorders in relation to distribution of hazardous air pollutants in the San Francisco Bay area. *Environmental Health Perspectives*, 114(9), 1438–1444.
- Wing, L. (1980). Childhood autism and social class: A question of selection? *British Journal of Psychiatry*, 137, 410–417.
- World Health Organisation (1993). *International statistical classification of diseases and related health problems (ICD-10)* (10th ed.). Geneva: World Health Organization.