Articles

Gene–environment correlations and causal effects of childhood maltreatment on physical and mental health: a genetically informed approach

Varun Warrier, Alex S F Kwong*, Mannan Luo*, Shareefa Dalvie, Jazz Croft, Hannah M Sallis, Jessie Baldwin, Marcus R Munafò, Caroline M Nievergelt, Andrew J Grant, Stephen Burgess, Tyler M Moore, Ran Barzilay, Andrew McIntosh, Marinus H van IJzendoorn, Charlotte A M Cecil

Summary

Background Childhood maltreatment is associated with poor mental and physical health. However, the mechanisms of gene–environment correlations and the potential causal effects of childhood maltreatment on health are unknown. Using genetics, we aimed to delineate the sources of gene–environment correlation for childhood maltreatment and the causal relationship between childhood maltreatment and health.

Methods We did a genome-wide association study meta-analysis of childhood maltreatment using data from the UK Biobank (n=143473), Psychiatric Genomics Consortium (n=26290), Avon Longitudinal Study of Parents and Children (n=8346), Adolescent Brain Cognitive Development Study (n=5400), and Generation R (n=1905). We included individuals who had phenotypic and genetic data available. We investigated single nucleotide polymorphism heritability and genetic correlations among different subtypes, operationalisations, and reports of childhood maltreatment. Family-based and population-based polygenic score analyses were done to elucidate gene–environment correlation mechanisms. We used genetic correlation and Mendelian randomisation analyses to identify shared genetics and test causal relationships between childhood maltreatment and mental and physical health conditions.

Findings Our meta-analysis of genome-wide association studies (N=185414) identified 14 independent loci associated with childhood maltreatment (13 novel). We identified high genetic overlap (genetic correlations 0.24-1.00) among different maltreatment operationalisations, subtypes, and reporting methods. Within-family analyses provided some support for active and reactive gene–environment correlation but did not show the absence of passive gene–environment correlation. Robust Mendelian randomisation suggested a potential causal role of childhood maltreatment in depression (unidirectional), as well as both schizophrenia and ADHD (bidirectional), but not in physical health conditions (coronary artery disease, type 2 diabetes) or inflammation (C-reactive protein concentration).

Interpretation Childhood maltreatment has a heritable component, with substantial genetic correlations among different operationalisations, subtypes, and retrospective and prospective reports of childhood maltreatment. Familybased analyses point to a role of active and reactive gene–environment correlation, with equivocal support for passive correlation. Mendelian randomisation supports a (primarily bidirectional) causal role of childhood maltreatment on mental health, but not on physical health conditions. Our study identifies research avenues to inform the prevention of childhood maltreatment and its long-term effects.

Funding Wellcome Trust, UK Medical Research Council, Horizon 2020, National Institute of Mental Health, and National Institute for Health Research Biomedical Research Centre.

Copyright © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Childhood maltreatment, including abuse and neglect, is a complex global problem, affecting up to 36% of the population worldwide.¹² According to most definitions, childhood maltreatment is inflicted by an agent who can be held responsible for the wellbeing of the child, with up to 80% of perpetrators being the parents or other family members of the child.²³

Parental experience of childhood maltreatment is an important risk factor for maltreating their own children.⁴⁵ This intergenerational transmission of childhood maltreatment might partly be due to shared social factors or to shared genetics between parents and their offspring. Studies that quantify the variance in childhood maltreatment attributable to genetics (broad-sense heritability), show childhood maltreatment has a twin and familial heritability of 6–62%.⁶⁻⁸ It is also possible to investigate heritability attributable to the additive effects of single nucleotide polymorphisms (SNPs), and a study from 2020 has shown a SNP heritability of 6% for childhood maltreatment.⁹

The heritable component of childhood maltreatment is not immutable and is thought to manifest through gene–environment correlations: passive (parental genes influence family environments and are inherited by their children), active (children's genes shape their behaviours





Lancet Psychiatry 2021; 8: 373–86

Published Online March 16, 2021 https://doi.org/10.1016/ S2215-0366(20)30569-1

See Comment page 348

*Contributed equally Department of Psychiatry (V Warrier PhD), **MRC Biostatistics Unit** (A J Grant PhD, S Burgess PhD), and Department of Public Health and Primary Care (S Burgess), University of Cambridge, Cambridge, UK; MRC Integrative Epidemiology Unit at the University of Bristol (A S F Kwong PhD, H M Sallis PhD, Prof M R Munafò PhD), **Department of Population** Health Sciences, Bristol Medical School (A S F Kwong, H M Sallis, (Croft PhD), and School of **Psychological Science** (Prof M R Munafò), University of Bristol, Bristol, UK; Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburah, UK (A S F Kwong): Department of Psychology, Education and Child Studies, Erasmus University Rotterdam Rotterdam Netherlands (M Luo Msc, Prof M H van Ilzendoorn PhD): Generation R Study Group. Erasmus MC. University Medical Center Rotterdam, Rotterdam, Netherlands (M Luo); South Africa MRC Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry and Neuroscience Institute, University of Cape Town, Cape Town, South Africa (S Dalvie PhD); Department of Clinical. Educational and Health Psychology, Division of Psychology and Language Sciences, University College London, London, UK (J Baldwin PhD, Prof M H van Ilzendoorn): Social.

Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK (I Baldwin): NIHR Biomedical **Research Centre at the** University Hospitals Bristol NHS Foundation Trust and the University of Bristol, Bristol, UK (Prof M R Munafò); Department of Psychiatry, University of California San Diego, La Jolla, CA, USA (Prof C M Nievergelt PhD); Center of Excellence for Stress and Mental Health (Prof C M Nievergelt) and Research Service (Prof C M Nievergelt), Veterans Affairs San Diego Healthcare System, San Diego, CA, USA (Prof C M Nievergelt): Department of Psychiatry, University of Pennsylvania, Philadelphia, PA, USA (T M Moore PhD, R Barzilav MD): Lifespan Brain Institute of the Children's Hospital of Philadelphia and University of Pennsylvania, Philadelphia, PA, USA (T M Moore, R Barzilay); Division of Psychiatry, Centre for Clinical Brain Sciences. (Prof A McIntosh MD) and Centre for Cognitive Ageing and Cognitive Epidemiology, School of Philosophy, Psychology and Language Sciences (Prof A McIntosh), University of Edinburgh, Edinburgh, UK; Department of Child and Adolescent Psychiatry (CAM Cecil PhD) and Department of Epidemiology (C A M Cecil), Erasmus MC. Rotterdam. Netherlands: Molecular Epidemiology, Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, Netherlands (C A M Cecil)

Correspondence to: Dr Varun Warrier, Department of Psychiatry, University of Cambridge, Cambridge CB2 8AH, UK vw260@medschl.cam.ac.uk

Research in context

Evidence before this study

Childhood maltreatment is known to be partly heritable but how different mechanisms of gene-environment correlation contribute to childhood maltreatment, and whether childhood maltreatment is causally related to mental and physical health outcomes is unknown. We searched PubMed and Google Scholar for genome-wide association studies and Mendelian randomisation analyses of childhood maltreatment from Jan 1, 1990, to Oct 1, 2020. Search terms were "childhood maltreatment" OR "childhood trauma" AND "GWAS" OR "genome-wide association study" OR "Mendelian Randomization", restricting the search to articles published in English. We found one genome-wide association study of childhood maltreatment with two identified genetic loci. We did not identify any studies investigating mechanisms of gene-environment correlations or causal effects of childhood maltreatment using Mendelian randomisation.

Added value of this study

In this genome-wide meta-analysis (N=185414) of childhood maltreatment—the largest to date to our knowledge we identified 14 independent genetic variants (13 novel) associated with childhood maltreatment. Using within-family polygenic score analyses, we found that this genetic signal is due to multiple gene–environment correlation mechanisms. We found some direct evidence for active and reactive gene–environment correlation, but could not show an absence

including risk-taking behaviour), and reactive or evocative (in which children's genes shape their behaviour and physical features, eliciting different responses from their parents or others).6,10-13 These mechanisms might vary among different groups of individuals. For example, children with ADHD and autism might be at an increased risk for childhood maltreatment compared to typically developing children,14,15 owing to mismatch between parental expectation and child behaviour (eg, parental expectation of greater social responsiveness and lower activity levels), which is a form of reactive geneenvironment correlation. Neither active or reactive geneenvironment correlation indicates that the child is to blame, as the responsibility for protecting a child lies primarily with the parents and additionally with society at large. No study, to our knowledge, has delineated the proportion of variance in childhood maltreatment explained by different gene-environment correlation mechanisms, which could help guide family-based prevention strategies.

Observational studies link childhood maltreatment with severe and long-lasting mental (eg, psychosis,¹⁶ depression,¹⁷ ADHD¹⁴) and physical health problems^{2,5} (eg, diabetes,¹⁸ cardiovascular disease¹⁹). Yet, there is considerable heterogeneity in these observations, partly because studies differ in how childhood maltreatment of passive gene-environment correlation. Finally, in this first Mendelian randomisation analyses of childhood maltreatment, to our knowledge, we identified potentially causal effects of childhood maltreatment on depression. We also identified bidirectional potentially causal effects between childhood maltreatment and both ADHD and schizophrenia. However, we found no evidence for causal effects of childhood maltreatment on physical health or inflammatory markers, or vice versa.

Implications of all the available evidence

Our study supports findings from observational epidemiological studies, providing the first evidence using Mendelian randomisation of potential causal relationships between childhood maltreatment and mental health conditions. These findings underscore the need to prevent childhood maltreatment and to identify mediating and moderating factors that can help minimise the effect of childhood maltreatment on mental health. However, in contrast to observational studies, we found no association between childhood maltreatment and physical health. Although childhood maltreatment is only partly heritable, delineating the source of the heritability is important as different gene-environment correlation mechanisms might indicate the need for different intervention strategies. Further research is needed to establish whether these gene-environment correlation mechanisms differ between populations and subtypes of childhood maltreatment.

is operationalised,²⁰ whether they examine the effects of specific subtypes,²¹ and whether prospective or retrospective reports of childhood maltreatment are used, which overlap only modestly.²² Furthermore, it is unclear whether observational associations are due to causal effects of childhood maltreatment, or to genetic or environmental confounding (eg, deprivation).^{25,23} For example, poor parental mental health is a risk factor for childhood maltreatment. Propensity for mental illness in children who are maltreated might be inherited from parents independently of having experienced childhood maltreatment (genetic confounding). Similarly, social deprivation is a risk for maltreatment and might also be independently associated with adverse health outcomes, thus presenting incorrectly as an association between adverse health and childhood maltreatment (environmental confounding). Identifying causal effects could help to mitigate adverse sequalae of childhood maltreatment.

Some studies have sought to investigate causality using co-twin control methods.^{24,25} However, ascribing causality in these studies is difficult, as twin studies rely on certain assumptions (eg, equal environment assumptions) and cannot exclude reverse causation, especially if longitudinal data are unavailable. It might be challenging to control for non-shared environments, particularly those that influence later health outcomes (eg, lifestyle choices and coronary artery disease). Mendelian randomisation²⁶ assumes that a genetic variant associated with an exposure can be used as an unconfounded proxy for that exposure to investigate its effect on potential outcomes. For example, a Mendelian randomisation study found a protective effect of exercise on major depressive disorder.²⁷ No study, to our knowledge, has used Mendelian randomisation to investigate the causal mechanisms of childhood maltreatment on mental and physical health.

Here, we present the largest genome-wide association study (GWAS) meta-analysis of childhood maltreatment to date, to our knowledge, to investigate the underlying mechanisms of childhood maltreatment, focusing on offspring DNA. We aimed to quantify the SNP heritability and genetic correlations across different operationalisations, subtypes, and reporting methods of childhood maltreatment. Finally, we investigated the genetic correlations and causal effects between childhood maltreatment and outcomes related to mental and physical health.

Methods

Overview

In this GWAS meta-analysis, we had access to individuallevel data from 159124 participants from four datasets: 143473 participants from the UK Biobank (birth year 1936–70);²⁸ 5400 from the Adolescent Brain Cognitive Development Study (birth year 2006–08);²⁹ 8346 from the Avon Longitudinal Study of Parents and Children (ALSPAC; birth year 1991–92);^{30,31} and 1905 from the Generation R Study (birth year 2002–06).^{32,33} In addition, we obtained summary GWAS statistics for the Psychiatric Genomics Consortium (PGC_26K) dataset (n=26 290 from 18 cohorts⁹). All participants (N=185414) were primarily of European genetic ancestry and provided information on childhood maltreatment. Further details are shown in the appendix (pp 5–7, 61–63).

Ethical approval was obtained from the Human Biology Research Ethics Committee, University of Cambridge (Cambridge, UK), and from the local cohort ethics committees.

Phenotypes

We define childhood maltreatment here as consisting of emotional, sexual, and physical abuse, and emotional and physical neglect. All retrospective reports of childhood maltreatment were self-reported, whereas most prospective reports of childhood maltreatment were reported by a parent or caregiver.

Participants in the UK Biobank completed the retrospectively reported five-item Childhood Trauma Screener.³⁴ This assessment consists of one question for each of the five trauma subtypes, each ranging from 0 (never true) to 4 (very often true), with total scores ranging from 0–20. Participants in the PGC_26K completed different retrospective questionnaires on childhood maltreatment, which included questions only

on sexual, physical, and emotional abuse. Total scores differed between the questionnaires, but all were coded continuously. In the Adolescent Brain Cognitive Development Study, parent-completed information on prospectively reported childhood maltreatment was used. This assessment comprised 13 questions on childhood maltreatment from the Kiddie Schedule for Affective Disorders and Schizophrenia-PTSD³⁵ and the Children's Report of Parental Behavior Inventory,36 with scores ranging from 0–13. We used continuous scores that were rank-based inverse normal transformed for the analyses. In ALSPAC, childhood maltreatment was prospectively recorded using multiple questionnaires at multiple instances (majority parent-report, several self-report), detailed elsewhere.¹⁶ Scores were binarised with any instance of childhood maltreatment indicated as 1 and no report of childhood maltreatment indicated as 0, in line with previous analyses.16 In Generation R, childhood maltreatment was prospectively measured using mothercompleted questionnaires including items on the Life Event and Difficulty Schedule,37 with scores ranging from 0-2, which were continuously coded. Further details are provided in the appendix (pp 7–10, 27, 61–63).

With the UK Biobank data, we investigated heritability of and genetic correlations among different subtypes and operationalisations of childhood maltreatment (appendix pp 7–10). For subtype analyses, we binarised the five phenotypes (sexual, emotional, and physical abuse, and emotional and physical neglect) due to skewness in the phenotypes. We defined four operationalisations of childhood maltreatment: a log-transformed sum-score of childhood maltreatment; a binary maltreatment score (0 *vs* any); a binary severe maltreatment score, in which scores 2–4 for each individual item were recorded as 1; and a binary severe childhood abuse score, which is the same as above but restricted to the three abuse items only.

GWAS meta-analysis and functional annotation

For each GWAS, age and sex were included as covariates. The primary GWAS in the UK Biobank was done using the BOLT-LMM, version 2.3.4, algorithm,38 with batch included as an additional covariate. GWAS in ALSPAC was done using BOLT-LMM, version 2.3.4,38 GWAS in Adolescent Brain Cognitive Development Study was done using FastGWA (using GCTA version 1.93.2),39 and we used linear regression using Plink,40 version 2.0, for Generation R. The linear mixed-effects models used in the Adolescent Brain Cognitive Development Study, UK Biobank, and ALSPAC account for both population stratification and relatedness. However, we included genetic principal components (up to 20) to accelerate the model identification process.³⁸ In Generation R, we included five genetic principal components as covariates to account for population stratification (appendix pp 10–13). We confirmed that there was no evidence of inflation in the test statistics of the GWAS by using the linkage disequilibrium score regression (LDSC)-based intercept.41 Sample-size weighted meta-analysis was done in METAL, version 2011-03-25.⁴² We did three meta-analyses. First, we meta-analysed the UK Biobank and the PGC_26K datasets to obtain a GWAS of retrospectively reported childhood maltreatment (GWAS_{retrospective}). Next, we meta-analysed Adolescent Brain Cognitive Development Study, ALSPAC, and Generation R datasets to obtain a GWAS of prospectively reported childhood maltreatment (GWAS_{prospective}). Finally, we meta-analysed all five datasets to obtain a GWAS of childhood maltreatment (GWAS_{childhoodmaltreatment}).

Independent significant loci were identified at a GWAS threshold of $p<5\times10^{-8}$, after clumping ($r^2=0.1$, 1000 kb), using the linkage disequilibrium weights generated from the European subset of the 1000 Genomes phase 3 dataset⁴³ in Plink, version 1.9. Functional annotation of top loci was done using expression quantitative trait loci data from GTEx,⁴⁴ BRAINEAC,⁴⁵ CommonMind Consortium,⁴⁶ and PsychEncode,⁴⁷ Hi-C data, positional mapping, and associations with other health-related phenotypes all using FUMA.⁴⁸

Gene identification was done using MAGMA,49 with the summary GWAS statistics. We identified significantly associated genes after Bonferroni correction. Enrichment for tissues and cell types was done using MAGMA and LDSC-specifically expressed genes (SEG),50 using summary GWAS statistics (appendix p 14). Specifically, using LDSC-SEG, we investigated enrichment for tissuespecific chromatin marks (ENCODE51 and Roadmap Epigenomics Project⁵²) and gene expression (GTEx), and corrected each of these analyses for multiple testing using Benjamini-Hochberg false discovery rate correction to account for the correlated nature of the variables tested and the enrichment for gene expression using Bonferroni correction. Using MAGMA, we investigated enrichment for genes with tissue-type (GTEx⁴⁴) and cell-type specific expression in neuronal cell types (PsychEncode⁴⁷), and corrected each of these analyses for multiple testing using Bonferroni correction.

SNP heritability and genetic correlations

SNP heritability of and genetic correlations between operationalisations and subtypes of childhood maltreatment were done using GCTA-GREML, version $1.93.2^{33}$ in a random subset of 19559 unrelated individuals from the UK Biobank (grm-cutoff for relatedness=0.05). For all analyses, we included year of birth, sex, genotyping batch, and the first 20 genetic principal components as covariates. Results were corrected for multiple testing using Benjamini-Hochberg false discovery rate correction owing to the correlated nature of the phenotypes.

Heritability analyses of the meta-analysed GWAS and other genetic correlations were done using LDSC⁵⁴ (appendix p 13), focusing on mental and physical health conditions, and psychological, behavioural, and anthropometric traits. We corrected for the 97 phenotypes tested using Bonferroni correction.

Polygenic score analyses

We investigated the variance explained by polygenic score (PGS) in two cohorts-a hold-out sample from the UK Biobank and ALSPAC. PGSs were calculated using PRSice-2⁵⁵ (clump $r^2=0.1$, 250kb, appendix pp 15–19). Only autosomes were included in the calculation of PGS as there is no consensus for how to handle sex chromosomes in PGS analyses.⁵⁶ To identify variance explained by the GWAS of childhood maltreatment, we did PGS analyses using GWAS_{retrospective} (base sample) in a hold-out sample of 12855 individuals from the UK Biobank (target sample). This process was done by doing a second GWAS of childhood maltreatment in the UK Biobank excluding the hold-out sample (log-transformed sum-score, n=130618) and meta-analysing the results with the PGC_26K. PGSs were generated in 9924 unrelated individuals from this subset of 12855 individuals at 11 p value thresholds. We included birth year, sex, genotyping batch, and the first 20 genetic principal components as covariates, and additionally Townsend Deprivation Index in a second model. We corrected for multiple testing using Benjamini-Hochberg false discovery rate correction.

In ALSPAC (target sample), we regressed the PGS for GWAS_{retrospective} (base sample) against binarised childhood maltreatment (prospectively reported) at four age groups (0-17, 0-4.9, 5-10.9, 11-17 years), with age (in months), sex, and the first ten genetic principal components as ALSPAC (maximum N=7453). We used only ten principal components as ALSPAC is a geographically-and subsequently, genetically-more homogenous cohort than the UK Biobank, and previous research has found poor evidence for population stratification in ALSPAC.⁵⁷ We restricted the PGS analyses in ALSPAC to PGS at a p value threshold of 1 as scores calculated at this threshold explained the highest variance in the hold-out sample from the UK Biobank. We corrected for the four different timepoints using Benjamini-Hochberg false discovery rate correction.

Contribution of different mechanisms to childhood maltreatment

We used three methods to delineate the contribution of different gene–environment correlation mechanisms to childhood maltreatment: (1) comparing between-sibling and between-family effects; (2) polygenic transmission disequilibrium tests in two autism cohorts; and (3) investigating the variance explained by PGS for retrospective childhood maltreatment in ALSPAC after accounting for well known familial risk factors for childhood maltreatment.

To quantify the variance explained by passive geneenvironment correlation and by active and reactive gene-environment correlation combined, we simultaneously investigated between-sibling and between-family effects of PGS (base sample $\text{GWAS}_{\text{childhoodmaltreatment}}^{58,59}$ in a hold-out sample of 12855 individuals from the UK Biobank (including 2849 sibling pairs, target sample; appendix pp 15–19), using a mixed-effects regression model with the following equation:

 $\begin{aligned} Childhood_maltreatment_{ij} &\sim \beta_{bsib}(PGS_{ij} - \overline{PGS}_j) \\ &+ \beta_{bfam}(\overline{PGS}_j) + Z_{(1..n)covariates} \end{aligned}$

where β_{bsib} is the between-sibling effect of PGS (representing reactive and active gene–environment correlation combined); β_{bjan} is the between-family effects; and (PGS_j) is the family-mean PGS. Covariates included were age, sex, genotyping batch, and 20 genetic principal components. Passive gene–environment correlation is estimated from the difference between β_{bsib} and β_{bjan} . SE was calculated by 10000 bootstraps. In the UK Biobank, the family mean is the sibling mean.

Although between-sibling PGS analysis assumes that a proportion of the familial environment is shared between siblings, this assumption might not always be true. An example is siblings who are discordant for neurodevelopmental conditions such as autism or ADHD. Because of the different support needs of the siblings, parental response to the two siblings will be different. This difference in familial environment between siblings indexes reactive gene-environment correlation, and to an extent active gene-environment correlation. To quantify this difference in gene-environment correlation, we did polygenic transmission disequilibrium tests60 in two cohorts: Simons Simplex Collection61 (n=2234 autistic individuals and n=1829 non-autistic siblings) and SPARK62 (n=2957 autistic individuals and n=1567 non-autistic siblings; appendix p 19) to investigate over-transmission of PGS for childhood maltreatment to autistic individuals versus non-autistic siblings.15,60 We use the term autistic as identity-first language is preferred by many autistic individuals.

Finally, we repeated the PGS analyses in ALSPAC, as outlined earlier, after including in separate models four parental risk factors5 of childhood maltreatment (ie, smoking, alcohol consumption, depression, and parental maltreatment). The risk factors were all measured prenatally to minimise the influence of a child's behaviour on parental phenotypes, leading to reactive gene-environment correlation (N=5988 to N=4508; see appendix p 70 for individual sample sizes). We corrected for multiple testing using Benjamini-Hochberg false discovery rate correction for each of the risk factors tested. Incomplete attenuation of the effects of PGS after accounting for known parental risk factors provided support for the active and reactive gene-environment correlation. We adjusted for eight phenotypes: parental depression assessed in the first trimester of pregnancy; parental depressive symptoms assessed in the second trimester; maternal depressive symptoms assessed in the third trimester and paternal depressive symptoms assessed in the second trimester; parental alcohol consumption assessed in the second trimester; parental alcohol consumption assessed in the third trimester; parental smoking assessed in the second trimester; parental smoking assessed in the third trimester; and parental history of childhood maltreatment, assessed across all trimesters.

Mendelian randomisation

We did two-sample, bidirectional Mendelian random isation analyses63,64 (appendix p 20) between childhood maltreatment and selected mental health outcomes (schizophrenia,65 major depressive disorder,66 bipolar disorder,67 ADHD,68 and autism69), physical health conditions (coronary artery disease⁷⁰ and type 2 diabetes⁷¹), and C-reactive protein as a marker of inflammation,72 and corrected with the Bonferroni correction. These phenotypes have all been associated with childhood maltreatment, and their GWAS does not include the UK Biobank, reducing bias in Mendelian randomisation estimates due to sample overlap.73 Bidirectional Mendelian randomisation was done using the following methods: inverse varianceweighted Mendelian randomisation, which assumes that all SNPs are valid instruments; median-weighted, which provides valid estimates even if up to 50% of the instruments are invalid;74 Mendelian randomisation-Egger, which accounts for pleiotropy by including an intercept term in the inverse variance-weighted model;75 and Mendelian randomisation-PRESSO, which accounts for pleiotropy by detecting and removing outliers.76 We additionally did leave-one-out analysis as a sensitivity check to investigate it the effects are driven by a subset of the variants.

Scripts and Summary GWAS statistics are available online.

Role of the funding source

The funders of the study had no role in study design, data analysis, data interpretation, or writing of the report.

Results

An overview of the study design is provided in figure 1. We did a series of analyses to quantify the heritabilities and genetic correlations of different operationalisations and subtypes of childhood maltreatment in the UK Biobank. Three of the four operationalisations had similar SNP heritability (0.093 [SE 0.019] to 0.056 [0.018], Δ SNP heritability all p values >0.05; figure 2A; see appendix p 63 for individual p values). However, restricting to severe abuse (binarised) identified lower SNP heritability (0.028 [SE 0.018]). All four operationalisations had modest to high genetic correlations with each other (genetic correlation (r_g) 0.47–1.00, figure 2B, appendix p 64), suggesting largely similar common variant genetics between the different operationalisations. We used log-transformed childhood maltreatment scores for

For **scripts** see https://github. com/vwarrier/ Childhoodmaltreatment_ genetics

For **summary statistics** see https://doi.org/10.17863/ CAM.65339



Figure 1: Study profile

PGC=Psychiatric Genomics Consortium. ALSPAC=Avon Longitudinal Study of Parents and Children. SNP=single nucleotide polymorphism. GWAS=genome-wide association study. pTDT=polygenic transmission disequilibrium test. SSC=Simons Simplex Collection.

subsequent analyses due to the relatively high SNP heritability compared with the other operationalisations of childhood maltreatment and high genetic correlations with other phenotypic operationalisations.

We next investigated the heritability and genetic correlations between subtypes of childhood maltreatment. We identified modest but significant SNP heritability for all five subtypes of childhood maltreatment (figure 2C, appendix p 65). Genetic correlations were high between childhood (overall) maltreatment and subtypes ($r_g \ 0.62-1.00$; figure 2D, appendix p 66). Genetic correlations among the five subtypes themselves ranged from low to high ($r_g \ 0.24-1.00$; figure 2D, appendix p 66).

Given the modest to high genetic correlations between operationalisations in the UK Biobank, we asked whether the two retrospective GWASs of childhood maltreatment were genetically correlated. We found a high genetic correlation between the UK Biobank and PGC_26K datasets ($r_g=0.64$ [SE=0.12]; p=1.15×10⁻⁷). Meta-analysis of the UK Biobank and PGC_26K (GWAS_{retrospective}) identified 257 significant SNPs representing 14 independent loci (figure 3, appendix pp 28, 67). In a hold-out sample of 9924 individuals from the UK Biobank, PGS from $GWAS_{retrospective}$ explained a maximum of 0.91% of the variance in retrospectively reported childhood maltreatment (p value threshold=1, $p=2\times10^{-16}$, appendix p 69). This result represents approximately 10% of the SNP heritability (0.093, or 9.3% of the total phenotypic variance). PGS at all 11 p value thresholds was significant after Benjamini-Hochberg false discovery rate correction. Accounting for Townsend deprivation index (a risk factor for childhood maltreatment)⁵ did not substantially attenuate the variance explained (p value threshold=1, r^2 =0.87%; p=2×10⁻¹⁶). Similarly, we found no change in SNP heritability of childhood maltreatment and subtypes after accounting for Townsend deprivation index (appendix p 64). Thus, the common variant signal for childhood maltreatment is independent of the heritable aspects of social deprivation.⁷⁷

To quantify the shared genetics between prospectively and retrospectively reported childhood maltreatment, we did a GWAS of prospectively reported childhood maltreatment (GWAS_{prospective} n=15650, SNP heritability 0.046 [SE 0.033]) by meta-analysing data from three datasets (Adolescent Brain Cognitive Development Study, ALSPAC, and Generation R). Genetic correlation between GWAS_{retrospective} and GWAS_{prospective} was high (r_g =0.72 [SE 0.36]; p=0.046), similar to that between the two GWAS of retrospectively reported childhood maltreatment (r_g =0.64 [SE 0.12]). Confirming this result, PGS for GWAS_{retrospective} explained a significant proportion of the variance in prospectively reported childhood maltreatment in ALSPAC across multiple age groups (Nagelkerke's pseudo r^2 =0.26%; p=2.04×10⁻⁷, appendix pp 29, 70–71).

Given the relatively high genetic correlation between retrospectively and prospectively reported childhood maltreatment, we did a GWAS meta-analysis of retrospectively and prospectively reported childhood maltreatment (GWAS_{childhoodmaltreatment}, Biobank. UK PGC_26K, ALSPAC, Adolescent Brain Cognitive Development Study, and Generation R). We identified 277 genome-wide significant SNPs, representing 14 independent loci (figure 3, appendix pp 30-44, 72). 12 of these lead SNPs had data available in $\text{GWAS}_{\text{prospective}}$. All showed a concordant effect direction between the $GWAS_{prospective}$ and $GWAS_{retrospective}$ (p=0.0005, binomial sign test), with six loci nominally significant (GWAS_{prospective} p<0.050) and one significant after Bonferroni correction for the 12 loci (rs3851357, GWAS_{prospective} p=0.0035). The meta-analysed GWAS of childhood maltreatment had a significant SNP heritability (0.079 [SE 0.0042], n=185414), and no inflation in statistics due to unresolved population stratification was detected (LDSC-based intercept 1.0076 [SE 0.0074]). Of the 14 loci, two were on the X-chromosome. Although our study was underpowered for sex-stratified GWAS, we note significant sex differences in overall childhood maltreatment and subtypes in the datasets in which we had access to individual level data. Notably, in most datasets, female individuals were more likely to report childhood maltreatment. In the UK Biobank, female individuals were more likely to report sexual abuse, while male individuals were more likely to report physical abuse in line with previous research (appendix pp 21-22).78

The genome-wide significant loci were significantly associated $(p<5\times10^{-8})$ with mental health problems



Figure 2: SNP heritability and genetic correlations between phenotypic operationalisations and subtypes of childhood maltreatment (A) SNP heritability of the four phenotypic definitions of childhood maltreatment and 95% CIs (n=19559). (B) Genetic correlations between the four phenotypic operationalisations of childhood maltreatment (n=19559). (C) SNP heritability of the subtypes of childhood maltreatment (binarised) alongside the log-transformed sum-score of childhood maltreatment and 95% CIs (n=19559). (D) Genetic correlations between subtypes of childhood maltreatment and log-transformed sum-score of childhood maltreatment (n=19559). SNP=single nucleotide polymorphism.

(six loci), risky behaviour (four loci), smoking and cannabis use (three loci), cardiovascular health, sleep difficulties, and reduced intelligence and educational attainment (two loci each) identified in previous GWASs (appendix pp 73–77).

Positional mapping identified 12 genes, expression quantitative trait loci identified 14 genes, chromatin interaction identified 15 genes (appendix pp 77–82), and MAGMA identified 19 significant genes after Bonferroni correction (appendix p 82). Four of these genes were identified by all four methods: *FES, FOXP2, SORCS3,* and *SAMD5.* The GWAS signal was significantly

enriched for foetal and adult brain-specific histone marks and DNAse hypersensitivity sites (appendix p 83), for genes with high expression in the excitatory neuron subpopulation (*Ex4*) in the adult post-mortem brain (appendix p 84), but not genes with tissue-specific expression (appendix pp 85–88).

To better understand the gene–environment correlation mechanisms that contribute to childhood maltreatment, we did three within-family PGS analyses. First, we compared between-sibling (active and reactive gene–environment correlation) with between-family (ie, total) PGS of GWAS_{childhoodmaltreatment} effects using linear



Figure 3: Circular Manhattan plot of childhood maltreatment Circular Manhattan plot for the GWAS meta-analysis of prospectively and retrospectively reported childhood maltreatment (n=185414, outer ring), and retrospectively reported childhood maltreatment (n=169766, inner ring). Red lines indicate GWAS significance threshold ($p=5 \times 10^{-8}$). The vertical axis provides the p values for single nucleotide polymorphisms included in the GWAS meta-analyses. GWAS=genome-wide association study.

mixed-effects models. The total between-family effect was significant (β =0.095 [SE 0.007]; p<2×10⁻¹⁶). We identified a small between-sibling effect (β =0.053 [0.02]; p=0.015), accounting for 58% of the total effect. By contrast, the difference in between-family and between-sibling PGS effects (passive gene–environment correlation) did not reach statistical significance (β =0.039 [SE 0.026]; p=0.13) accounting for approximately 42% of the total effect. The difference between the combined active and reactive gene–environment correlation versus passive gene–environment correlation versus passive gene–environment correlation did not reach statistical significance (Δ PGS=0.014 [SE 0.05]; p=0.77; figure 4A).

The between-sibling analyses assumed that parenting (and consequently, familial environment) was similar, or shared, between siblings. However, parenting could systematically differ between, for example, autistic individuals and their non-autistic siblings, in which case the proportion of shared familial environment might be smaller than for families without autistic children. In these families, a larger proportion of sibling difference in childhood maltreatment could be attributable to reactive (eg, poor understanding and support) and active gene-environment correlation (eg, greater risk-taking behaviour, lower social responsiveness) than for siblings with no known neurodevelopmental disorders. Supporting this hypothesis, we found an over-transmission of PGS of GWAS_{childhoodmaltreatment} to autistic children from their parents (mean difference 0.056 [SE 0.02]; p=0.010) but not to their non-autistic siblings (mean difference -0.01 [0.02]; p=0.56) in the Simons Simplex Collection, suggesting that an increased risk for childhood maltreatment in autistic individuals was partly explained by increased active and reactive gene–environment correlation. This finding was replicated in the SPARK dataset (autistic individuals mean difference 0.059 [SE 0.02]; p=1.1×10⁻³; non-autistic siblings mean difference 0.01 [0.025]; p=0.65; figure 4B).

We used the family data in ALSPAC to investigate whether accounting for four known parental risk factors (alcohol consumption, childhood maltreatment, depression, and smoking—measured prenatally at multiple timepoints) for childhood maltreatment substantially attenuated the variance explained by PGS for GWAS_{retrospective}. None of the four risk factors substantially reduced the variance explained by PGS (figure 4C; appendix pp 29, 71–72).

We next investigated the shared genetics between childhood maltreatment and several health-related phenotypes to contextualise the GWAS_{childhoodmaltreatment} signal. After Bonferroni correction, we identified significant positive genetic correlations between childhood maltreatment and mental health conditions (eg, schizophrenia and depression), anthropometric traits (eg, obesity and body-mass index), insomnia, and coronary artery disease. Childhood maltreatment also had modest negative genetic correlations with intelligence, educational attainment, and age of onset traits (eg, smoking, parental death, first birth; appendix pp 88–93).

To investigate the potential bidirectional causal effect between childhood maltreatment and selected mental and physical health conditions, we did Mendelian randomisation analyses. Inverse varianceweighted Mendelian randomisation analyses found a significant causal effect of childhood maltreatment on major depressive disorder (figure 5A) but not vice versa (figure 5D). We found a significant bidirectional causal effect for childhood maltreatment with ADHD and schizophrenia (figure 5, appendix p 93). Sensitivity analyses using the weighted median method and Mendelian randomisation-PRESSO were also significant between childhood maltreatment and depression, ADHD, and schizophrenia. Mendelian randomisation-Egger analyses were significant only for the bidirectional effect between childhood maltreatment and schizophrenia (appendix pp 45-60, 93). Leave-one-out analyses did not suggest that any one genetic variant was driving the result, despite the locus-level pleiotropy, and Steiger analyses indicated that the causal direction between the exposure and outcome was correct in all analyses. By contrast, we did not identify a significant causal effect of childhood maltreatment on autism, bipolar disorder, coronary artery disease, type 2 diabetes, C-reactive protein concentration, or vice versa (appendix pp 45-60, 93).

Discussion

In this study, we found modest SNP heritability and substantial genetic correlations among different operationalisations, subtypes, and reports of childhood maltreatment. This GWAS meta-analysis of childhood maltreatment in 185414 individuals identified 14 genetic variants (13 novel). Our findings provide some evidence for the role of active and reactive gene–environment correlation in childhood maltreatment, although passive gene–environment correlation could not be excluded. Mendelian randomisation analyses found a unidirectional causal effect of childhood maltreatment on depression, bidirectional effects with ADHD and schizophrenia, but no evidence for unidirectional or bidirectional causal effects on physical health problems.

Childhood maltreatment is modestly heritable, with SNP heritability similar to that of depression.79 This SNP heritability provides a lower-bound for the variance attributable to genetic effects. Overall, we identify modest to high genetic correlations among different subtypes, operationalisations, and reports of childhood maltreatment. A 2019 meta-analysis²² found low agreement between prospectively and retrospectively reported childhood maltreatment, accounting for chance overlap. However, the raw agreement identified by the same study was 76%. Although it is difficult to compare phenotypic overlap with genetic correlation directly, we believe genetic correlations are more similar to raw agreement as chance overlap is not accounted for in genetic correlation analyses. Furthermore, the proportion of variance attributable to SNPs is modest, suggesting that factors that contribute to relatively low overlap between prospective and retrospective reports might be non-genetic or due to genetic effects not tagged by the SNPs tested.

Genome-wide analyses identified 14 independent loci for childhood maltreatment. The GWAS signal was enriched for regulatory chromatin marks in brain tissues and for genes that are highly expressed in excitatory neurons. These loci are implicated in mental health disorders, risky behaviours, cardiovascular disease, and intelligence, suggesting considerable locus-level

Figure 4: Delineating different gene-environment correlation mechanisms (A) Regression estimates and 95% CIs of active and reactive, passive, and total effects of childhood maltreatment PGS on childhood maltreatment (n=12855 included 2849 sibling pairs). (B) Polygenic transmission disequilibrium test to investigate over-transmission of childhood maltreatment polygenic scores from parents to autistic children and non-autistic siblings in two cohorts: SSC (n=2234 autistic individuals and 1829 non-autistic siblings) and SPARK (n=2957 autistic individuals and 1567 non-autistic siblings). Mean PGS over-transmission (difference between standardised parental mean PGS and standardised child mean PGS) and 95% CIs provided. (C) Effect of childhood maltreatment PGS on prospectively measured childhood maltreatment (age 0-17 years) in Avon Longitudinal Study of Parents and Children (unadjusted model). The models were adjusted for parental alcohol consumption, parental depression, parental experience of childhood maltreatment, and parental smoking, all measured at the time of pregnancy. Odds ratios and 95% CIs are provided. Sample sizes are provided in the appendix (pp 71-72). PGS=polygenic scores. SSC=Simons Simplex Collection.





Figure 5: Mendelian randomisation analyses

Scatter plots of the SNP effects of childhood maltreatment on major depressive disorder (A), schizophrenia (B), and ADHD (C). Scatter plots of the SNP effects of major depressive disorder (D), schizophrenia (E), and ADHD (F) on childhood maltreatment. All units of associations are log-odds ratios. Slopes provided correspond to three different Mendelian randomisation methods used (inverse variance-weighted, weighted median, and Mendelian randomisation-Egger). The Mendelian randomisation-Egger intercept was significant only for the causal effect of childhood maltreatment on schizophrenia (B; p=0.012). SNP=single nucleotide polymorphism.

pleiotropy. Although we did not test this, the identified genomic variants might interact with exposure to childhood maltreatment to influence epigenetic patterns, thereby contributing to the risk of developing mental and physical health disorders.⁸⁰

Within-family analyses identified significant estimates for active and reactive gene–environment correlation (58%). However, there was no statistical evidence for a difference between passive gene–environment correlation and active and reactive gene–environment correlation. Providing additional evidence for the role of active and reactive gene–environment correlation, childhood maltreatment PGSs are over-transmitted in autistic individuals. It remains unclear which aspects of autism might explain such gene–environment correlation. Importantly, this analysis does not test for the presence or absence of passive gene–environment correlation as all analyses were within-family analyses. Finally, accounting for four known parental risk factors for childhood maltreatment (alcohol consumption, childhood maltreatment, parental depression, and smoking) did not substantially attenuate the variance explained by PGS in prospectively reported childhood maltreatment. These parental risk factors were, however, minimally phenotyped, and we did not have data to test for other parental risk factors (eg, aggression⁵). Together, the analyses support the combined role of active and reactive gene–environment correlation, but do not show the absence of passive gene–environment correlation. Better delineation of the mechanisms of gene– environment correlation can help improve family-based support, for instance, by directly addressing parental mental health in the case of passive gene–environment correlation, or coaching parents on positive discipline methods in the case of reactive and active gene– environment correlation. Further research in larger and more diverse samples is needed to inform interventions.

Given the complex antecedents and consequences of childhood maltreatment, we investigated genetic correlations between childhood maltreatment and several health-related traits. Childhood maltreatment was genetically correlated with some mental health, reproductive, and physical health traits. Notably, childhood maltreatment was negatively genetically correlated with a number of age-of-onset traits, in line with the life history theory-an evolutionary framework suggesting that childhood maltreatment leads to accelerated development and reproduction, with long-term costs for mental and physical health.⁸¹ However, these genetic correlations do not imply causality as the proposed associations could be due to genetic or environmental confounding or pleiotropy, all of which can be reflected in genetic correlation analyses.

Mendelian randomisation analyses found a potential causal effect of childhood maltreatment on major depressive disorder, and bidirectional causal effects on ADHD and schizophrenia, consistent with observational studies.14,16,17 The bidirectional causal effect could be explained by either passive gene-environment correlation (parents of children with ADHD and psychosis have elevated traits in similar domains and might maltreat their children,5 in turn contributing to the mental health problems) or by active and reactive geneenvironment correlation (PGSs for schizophrenia and ADHD are associated with childhood difficulties,57,82 leading to increased risk of maltreatment and subsequent mental health problems). In addition to existing literature using co-twin designs, this study provides evidence for the causal role of childhood maltreatment in some mental health disorders.^{24,25}

We did not identify potential causal effects of autism on childhood maltreatment, in contrast to the polygenic transmission disequilibrium test results. This finding might be because the weak instrument used in the autism Mendelian randomisation analysis consisted of only four SNPs or because of the underlying heterogeneity in autism.⁸³ The heterogeneity is unlikely to affect within-family studies, but would influence the GWAS hits identified. Finally, the observed polygenic transmission disequilibrium test results could arise if both childhood maltreatment and autism were downstream of a common causal factor, but neither caused the other, an association that need not be reflected in Mendelian randomisation analyses. The current analyses found no causal relationship between childhood maltreatment and physical health conditions, which might be due to the relatively weak instruments (variance in exposure explained by the instruments was low, appendix pp 45–60) or because the observed associations were correlational rather than causal.

There are limitations to this study. First, common genetic variants explain a relatively small proportion of the variance in childhood maltreatment, thus non-genetic factors cannot be ignored. Genetic correlations and geneenvironment correlation effects must be interpreted while keeping the modest SNP heritability in mind. For example, although there is a high genetic correlation between childhood maltreatment and some mental health conditions, the covariance will be low due to the modest SNP heritability. Second, maltreatment was not uniformly measured across cohorts, increasing variability due to non-harmonised data collection, including age when maltreatment was measured. Third, we do not distinguish between objective childhood maltreatment and subjective valence of childhood maltreatment.22 Fourth, although our study considers childhood maltreatment as a single phenotype (to increase statistical power) and we found considerable genetic correlation between operationalisations, subtypes, and reports of childhood maltreatment, there could be subtype-specific and sex-specific genetic effects. Fifth, individuals with severe mental health disorders are less likely to answer follow-up questionnaires in the UK Biobank,84 and the current GWAS reflects ascertainment bias. Finally, our analyses included only individuals of European ancestries due to limitations in data availability.

The disappointing efficacy of current interventions to curb child maltreatment5 requires new ideas about potential mechanisms to be targeted as well as biological pathways that might mediate the long-term effect of maltreatment. Our findings of a significant heritability do not imply that environmental factors are absent, that the child is to blame, or that the heritability is fixed. These ideas are discussed in the appendix (pp 93–95). Rather, our findings provide empirical support that the mechanisms underlying maltreatment are complex, reflecting in part the action of multiple forms of geneenvironment correlation. The correlation mechanisms investigated here probably contribute to intergenerational transmission of childhood maltreatment, as both immediate environment and genes are inherited by children from parents. These findings highlight the importance of family-based support strategies, targeting parents and their interaction with their children to minimise the risk of child maltreatment, and limiting intergenerational transmission of childhood maltreatment.

Contributors

VW, ASFK, ML, and SD were responsible for data analysis. VW, ASFK, ML, MHvI, and CAMC were responsible for study design. VW, ASFK,

ML, SD, JC, HMS, CMN, MHvI, and CAMC were responsible for data collection. VW, ASFK, ML, SD, JC, HMS, JB, MRM, CMN, AJG, SB, TM, RB, AM, MHvI, and CAMC were responsible for data interpretation. VW, ASFK, ML, SD, JC, HMS, JB, MRM, CMN, AJG, SB, TM, RB, AM, MHvI, and CAMC were responsible for writing. The corresponding author had full access to all the data, with the exception of data from ALSPAC and Generation R, and the final responsibility to submit for publication. ASFK and JC had access to all the data in ALSPAC and ML and CAMC had access to all the data in Generation R.

Declaration of interests

AMM has received research support from Eli Lilly, Janssen, Pfizer, and the Sackler Foundation. RB serves on the scientific board and reports stock ownership in Taliaz Health, with no conflict of interest relevant to this work. MHvI is a founder and member of the non-profit Beagle Advice, Research and Development company, with no conflict of interests with the current work. Some of his work on child maltreatment has been supported by the Lumos Foundation, a charity that financed some activities of *The Lancet* Commission on Institutionalisation and Deinstitutionalisation of Children, of which he was a member. All other authors declare no competing interests.

Data sharing

The study website contains details of all data available through a fully searchable data dictionary. Some of these data were collected using REDCap.

This work was supported by the Wellcome Trust (grant

number 214322/Z/18/Z, 104036/Z/14/Z, 204623/Z/16/Z,

Acknowledgments

For details of **REDCap** see https://projectredcap.org/ resources/citations/

www.bristol.ac.uk/alspac/

researchers/our-data/

For data dictionary see http://

and 217065/Z/19/Z). VW was funded by the Bowring Research Fellowship from St Catharine's College (Cambridge, UK), and the Wellcome Trust Collaborative Award (grant number 214322/Z/18/Z). ASFK and AM are supported by the Wellcome Trust (grant number 104036/Z/14/Z). ASFK is also supported by an ESRC Postdoctoral Fellowship (grant number ES/V011650/1). ML is supported by a scholarship from the China Scholarship Council (grant number 201706990036). CAMC has received funding from the EU's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie grant agreement (707404) and grant agreement (848158; EarlyCause Project). MHvI is supported by the Dutch Ministry of Education, Culture, and Science and the Netherlands Organization for Scientific Research (grant number 024.001.003, Consortium on Individual Development) and by a Spinoza Prize of the Netherlands Organization for Scientific Research. HMS and MRM are supported by the Medical Research Council and the University of Bristol (grant number MC_UU_00011/7) and by the National Institute for Health Research (NIHR) Biomedical Research Centre at the University Hospitals Bristol National Health Service Foundation Trust and the University of Bristol. HMS is also supported by the European Research Council (grant number 758813 MHINT). CMN is supported by the National Institute for Mental Health (NIMH; grant number R01MH106595) and the Center of Excellence for Stress and Mental Health, Veterans Affairs San Diego. AJG and SB are supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society (grant number 204623/Z/16/Z). TMM and RB are supported by the NIMH (grant number R01MH117014, TMM; K23MH120437, RB). The research was done in association with the NIHR Cambridge Biomedical Research Centre, and the NIHR Collaboration for Leadership in Applied Health Research and Care East of England at Cambridgeshire and Peterborough NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the National Health Service, the NIHR, or the Department of Health and Social Care. This research was possible due to two applications to the UK Biobank (projects 20904 and 23787). This research was co-funded by the NIHR Cambridge Biomedical Research Centre and a Marmaduke Sheild grant. The UK Medical Research Council and the Wellcome Trust (grant number 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. A comprehensive list of grant funding is available on the ALSPAC website. We are very grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers,

For **grant funding** see http:// www.bristol.ac.uk/alspac/ external/documents/grantacknowledgements.pdf receptionists, and nurses. The first phase of the Generation R Study was made possible by financial support from the Erasmus Medical Centre, Rotterdam; the Erasmus University Rotterdam; ZonMw; the Netherlands Organization for Scientific Research; and the Ministry of Health, Welfare and Sport. The authors gratefully acknowledge the contribution of all children and parents, general practitioners, hospitals, midwives, and pharmacies involved in the Generation R Study. The Generation R study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Erasmus School of Social and Behavioural Sciences at Erasmus University Rotterdam; the Municipal Health Service Rotterdam area, Rotterdam; the Rotterdam Homecare Foundation, Rotterdam; and the Stichting Trombosedienst and Artsenlaboratorium Rijnmond, Rotterdam. We thank Veera M Rajagopal for helpful discussions. We thank Saskia Selzam for providing us with scripts to analyse some of the data.

References

- Stoltenborgh M, Bakermans-Kranenburg MJ, Alink LRA, van IJzendoorn MH. The prevalence of child maltreatment across the globe: review of a series of meta-analyses. *Child Abuse Rev* 2015; **24**: 37–50.
- 2 Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet* 2009; 373: 68–81.
- 3 Hughes K, Bellis MA, Hardcastle KA, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health* 2017; 2: e356–66.
- 4 Van Wert M, Anreiter I, Fallon BA, Sokolowski MB. Intergenerational transmission of child abuse and neglect: a transdisciplinary analysis. *Gend Genome* 2019; 3: 247028971982610.
- 5 van IJzendoorn MH, Bakermans-Kranenburg MJ, Coughlan B, Reijman S. Annual research review: umbrella synthesis of metaanalyses on child maltreatment antecedents and interventions: differential susceptibility perspective on risk and resilience. J Child Psychol Psychiatry 2020; 61: 272–90.
- 6 Pezzoli P, Antfolk J, Hatoum AS, Santtila P. Genetic vulnerability to experiencing child maltreatment. *Front Genet* 2019; 10: 852.
- 7 Fisher HL, Caspi A, Moffitt TE, et al. Measuring adolescents' exposure to victimization: the environmental risk (E-Risk) longitudinal twin study. *Dev Psychopathol* 2015; 27: 1399–416.
- 8 Pittner K, Bakermans-Kranenburg MJ, Alink LRA, et al. Estimating the heritability of experiencing child maltreatment in an extended family design. *Child Maltreat* 2020; 25: 289–99.
- 9 Dalvie S, Maihofer AX, Coleman JRI, et al. Genomic influences on self-reported childhood maltreatment. *Transl Psychiatry* 2020; 10: 38.
- 10 Knafo A, Jaffee SR. Gene-environment correlation in developmental psychopathology. Dev Psychopathol 2013; 25: 1–6.
- 11 Jaffee SR, Price TS. Gene-environment correlations: a review of the evidence and implications for prevention of mental illness. *Mol Psychiatry* 2007; 12: 432–42.
- 12 Plomin R, DeFries JC, Loehlin JC. Genotype-environment interaction and correlation in the analysis of human behavior. *Psychol Bull* 1977; 84: 309–22.
- 13 Griffiths LJ, Wolke D, Page AS, Horwood JP. Obesity and bullying: different effects for boys and girls. Arch Dis Child 2006; 91: 121–25.
- 14 Sanderud K, Murphy S, Elklit A. Child maltreatment and ADHD symptoms in a sample of young adults. *Eur J Psychotraumatol* 2016; 7: 32061.
- 15 Warrier V, Baron-Cohen S. Childhood trauma, life-time self-harm, and suicidal behaviour and ideation are associated with polygenic scores for autism. *Mol Psychiatry* 2019; published online Oct 29. https://doi.org/10.1038/s41380-019-0550-x.
- 16 Croft J, Heron J, Teufel C, et al. Association of trauma type, age of exposure, and frequency in childhood and adolescence with psychotic experiences in early adulthood. *JAMA Psychiatry* 2019; 76: 79–86.
- 17 Scott KM, Smith DR, Ellis PM. Prospectively ascertained child maltreatment and its association with DSM-IV mental disorders in young adults. Arch Gen Psychiatry 2010; 67: 712–19.
- 18 Shields ME, Hovdestad WE, Pelletier C, Dykxhoorn JL, O'Donnell SC, Tonmyr L. Childhood maltreatment as a risk factor for diabetes: findings from a population-based survey of Canadian adults. BMC Public Health 2016; 16: 879.

- 19 Suglia SF, Koenen KC, Boynton-Jarrett R, et al. Childhood and adolescent adversity and cardiometabolic outcomes: a scientific statement from the American Heart Association. *Circulation* 2018; 137: e15–28.
- 20 Jackson Y, McGuire A, Tunno AM, Makanui PK. A reasonably large review of operationalization in child maltreatment research: assessment approaches and sources of information in youth samples. *Child Abuse Negl* 2019; 87: 5–17.
- 21 Clemens V, Huber-Lang M, Plener PL, Brähler E, Brown RC, Fegert JM. Association of child maltreatment subtypes and longterm physical health in a German representative sample. *Eur J Psychotraumatol* 2018; 9: 1510278.
- 22 Baldwin JR, Reuben A, Newbury JB, Danese A. Agreement between prospective and retrospective measures of childhood maltreatment: a systematic review and meta-analysis. *JAMA Psychiatry* 2019; 76: 584–93.
- 23 South SC, Schafer MH, Ferraro KF. Genetic and environmental overlap between childhood maltreatment and adult physical health. *Twin Res Hum Genet* 2015; 18: 533–44.
- 24 Schaefer JD, Moffitt TE, Arseneault L, et al. Adolescent victimization and early-adult psychopathology: approaching causal inference using a longitudinal twin study to rule out noncausal explanations. *Clin Psychol Sci* 2018; **6**: 352–71.
- 25 Brown RC, Berenz EC, Aggen SH, et al. Trauma exposure and axis I psychopathology: a co-twin control analysis in Norwegian young adults. *Psychol Trauma* 2014; 6: 652–60.
- 26 Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet* 2014; 23: R89–98.
- 27 Choi KW, Chen CY, Stein MB, et al. Assessment of bidirectional relationships between physical activity and depression among adults a 2-sample Mendelian randomization study. *JAMA Psychiatry* 2019; **76**: 399–408.
- 28 Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* 2015; 12: e1001779.
- 29 Casey BJ, Cannonier T, Conley MI, et al. The Adolescent Brain Cognitive Development (ABCD) study: imaging acquisition across 21 sites. *Dev Cogn Neurosci* 2018; **32**: 43–54.
- 30 Fraser A, Macdonald-Wallis C, Tilling K, et al. Cohort profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol* 2013; 42: 97–110.
- 31 Boyd A, Golding J, Macleod J, et al. Cohort profile: the 'children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 2013; 42: 111–27.
- 32 Hofman A, Jaddoe VWV, Mackenbach JP, et al. Growth, development and health from early fetal life until young adulthood: the Generation R Study. *Paediatr Perinat Epidemiol* 2004; 18: 61–72.
- 33 Kooijman MN, Kruithof CJ, van Duijn CM, et al. The Generation R Study: design and cohort update 2017. Eur J Epidemiol 2016; 31: 1243–64.
- 34 Glaesmer H, Schulz A, Häuser W, Freyberger HJ, Brähler E, Grabe H-J. Der Childhood Trauma Screener (CTS) - Entwicklung und Validierung von Schwellenwerten zur Klassifikation. *Psychiatr Prax* 2013; 40: 220–26 (in German).
- 35 Geller B, Zimerman B, Williams M, et al. Reliability of the Washington University in St. Louis Kiddie Schedule for Affective Disorders and Schizophrenia (WASH-U-KSADS) mania and rapid cycling sections. J Am Acad Child Adolesc Psychiatry 2001; 40: 450–55.
- 36 Schaefer ES. Children's reports of parental behavior: an inventory. *Child Dev* 1965; **36**: 413–24.
- 37 Dunn EC, Nishimi K, Neumann A, et al. Time-dependent effects of exposure to physical and sexual violence on psychopathology symptoms in late childhood: in search of sensitive periods in development. J Am Acad Child Adolesc Psychiatry 2020; 59: 283–95.
- 38 Loh P-R, Kichaev G, Gazal S, Schoech AP, Price AL. Mixed-model association for biobank-scale datasets. Nat Genet 2018; 50: 906–08.
- 39 Jiang L, Zheng Z, Qi T, et al. A resource-efficient tool for mixed model association analysis of large-scale data. *Nat Genet* 2019; 51: 1749–55.
- 40 Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007; 81: 559–75.

- 41 Bulik-Sullivan BK, Loh P-R, Finucane HK, et al. LD score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015; 47: 291–95.
- 42 Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient metaanalysis of genomewide association scans. *Bioinformatics* 2010; 26: 2190–91.
- 43 Delaneau O, Marchini J, McVean GA, et al. Integrating sequence and array data to create an improved 1000 Genomes Project haplotype reference panel. *Nat Commun* 2014; 5: 3934.
- 44 Lonsdale J, Thomas J, Salvatore M, et al. The Genotype-Tissue Expression (GTEx) project. *Nat Genet* 2013; **45**: 580–85.
- 45 Ramasamy A, Trabzuni D, Guelfi S, et al. Genetic variability in the regulation of gene expression in ten regions of the human brain. *Nat Neurosci* 2014; 17: 1418–28.
- 46 Hoffman GE, Bendl J, Voloudakis G, et al. CommonMind Consortium provides transcriptomic and epigenomic data for schizophrenia and bipolar disorder. Sci Data 2019; 6: 180.
- 47 Akbarian S, Liu C, Knowles JA, et al. The PsychENCODE project. Nat Neurosci 2015; 18: 1707–12.
- 48 Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation of genetic associations with FUMA. *Nat Commun* 2017; 8: 1826.
- 49 de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput Biol* 2015; 11: e1004219.
- 50 Finucane HK, Reshef YA, Anttila V, et al. Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat Genet* 2018; 50: 621–29.
- 51 Bernstein BE, Birney E, Dunham I, Green ED, Gunter C, Snyder M. An integrated encyclopedia of DNA elements in the human genome. *Nature* 2012; 489: 57–74.
- 52 Kundaje A, Meuleman W, Ernst J, et al. Integrative analysis of 111 reference human epigenomes. *Nature* 2015; **518**: 317–30.
- 53 Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genomewide complex trait analysis. Am J Hum Genet 2011; 88: 76–82.
- 54 Bulik-Sullivan B, Finucane HK, Anttila V, et al. An atlas of genetic correlations across human diseases and traits. *Nat Genet* 2015; 47: 1236–41.
- 55 Choi SW, O'Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience* 2019; 8: giz082.
- 56 Choi SW, Mak TSH, O'Reilly PF. Tutorial: a guide to performing polygenic risk score analyses. *Nat Protoc* 2020; 15: 2759–72.
- 57 Riglin L, Collishaw S, Richards A, et al. Schizophrenia risk alleles and neurodevelopmental outcomes in childhood: a populationbased cohort study. *Lancet Psychiatry* 2017; 4: 57–62.
- 58 Selzam S, Ritchie SJ, Pingault JB, Reynolds CA, O'Reilly PF, Plomin R. Comparing within- and between-family polygenic score prediction. Am J Hum Genet 2019; 105: 351–63.
- 59 Demange PA, Hottenga JJ, Abdellaoui A, et al. Parental influences on offspring education: indirect genetic effects of non-cognitive skills. *bioRxiv* 2020; published online Sept 16. https://doi. org/10.1101/2020.09.15.296236 (preprint).
- 60 Weiner DJ, Wigdor EM, Ripke S, et al. Polygenic transmission disequilibrium confirms that common and rare variation act additively to create risk for autism spectrum disorders. *Nat Genet* 2017; 49: 978–85.
- 61 Fischbach GD, Lord C. The Simons Simplex Collection: a resource for identification of autism genetic risk factors. *Neuron* 2010; 68: 192–95.
- 62 Feliciano P, Daniels AM, Snyder LG, et al. SPARK: a US cohort of 50,000 families to accelerate autism research. *Neuron* 2018; 97: 488–93.
- 63 Slob EAW, Burgess S. A comparison of robust Mendelian randomization methods using summary data. *Genet Epidemiol* 2020; 44: 313–29.
- 64 Byrne EM, Yang J, Wray NR. Inference in psychiatry via 2-sample Mendelian randomization-from association to causal pathway? JAMA Psychiatry 2017; 74: 1191–92.
- 65 Lam M, Chen C-Y, Li Z, et al. Comparative genetic architectures of schizophrenia in East Asian and European populations. *Nat Genet* 2019; 51: 1670–78.
- 66 Wray NR, Ripke S, Mattheisen M, et al. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet* 2018; 50: 668–81.

- 67 Stahl EA, Breen G, Forstner AJ, et al. Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat Genet* 2019; 51: 793–803.
- 68 Demontis D, Walters RK, Martin J, et al. Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet* 2019; 51: 63–75.
- 69 Grove J, Ripke S, Als TD, et al. Identification of common genetic risk variants for autism spectrum disorder. *Nat Genet* 2019; 51: 431–44.
- 70 Nikpay M, Goel A, Won HH, et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet* 2015; 47: 1121–30.
- 71 Morris AP, Voight BF, Teslovich TM, et al. Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat Genet* 2012; 44: 981–90.
- 72 Wojcik GL, Graff M, Nishimura KK, et al. Genetic analyses of diverse populations improves discovery for complex traits. *Nature* 2019; 570: 514–18.
- 73 Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-sample Mendelian randomization. *Genet Epidemiol* 2016; 40: 597–608.
- 74 Bowden J, Davey Smith G, Haycock PC, Burgess S. Consistent estimation in Mendelian randomization with some invalid instruments using a weighted median estimator. *Genet Epidemiol* 2016; 40: 304–14.
- 75 Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015; 44: 512–25.
- 76 Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018; 50: 693–98.

- 77 Hill WD, Hagenaars SP, Marioni RE, et al. Molecular genetic contributions to social deprivation and household income in UK Biobank. *Curr Biol* 2016; 26: 3083–89.
- 78 Sunley AK, Lok A, White MJ, et al. Ethnic and sex differences in the association of child maltreatment and depressed mood. The HELIUS study. *Child Abuse Negl* 2020; **99**: 104239.
- 79 Howard DM, Adams MJ, Clarke TK, et al. Genome-wide metaanalysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci* 2019; 22: 343–52.
- 80 Cecil CAM, Zhang Y, Nolte T. Childhood maltreatment and DNA methylation: a systematic review. *Neurosci Biobehav Rev* 2020; 112: 392–409.
- 81 Cabeza de Baca T, Ellis BJ. Early stress, parental motivation, and reproductive decision-making: applications of life history theory to parental behavior. *Curr Opin Psychol* 2017; 15: 1–6.
- 82 Hamshere ML, Langley K, Martin J, et al. High loading of polygenic risk for ADHD in children with comorbid aggression. *Am J Psychiatry* 2013; **170**: 909–16.
- 83 Warrier V, Toro R, Won H, et al. Social and non-social autism symptoms and trait domains are genetically dissociable. *Commun Biol* 2019; 2: 328.
- 84 Adams MJ, Hill WD, Howard DM, et al. Factors associated with sharing e-mail information and mental health survey participation in large population cohorts. *Int J Epidemiol* 2019; 49: 410–21.